


CAT(0) spaces

A geodesic metric space X is CAT(0) if for every geodesic triangle $P = \Delta(p, q, r) \subseteq X$ there exists a comparison triangle in \mathbb{E}^2 with the same side lengths as P such that for each pair of points $x, y \in \partial P$ we have:



A group is CAT(0) if it acts properly cocompactly by isometries on a CAT(0) space.

Examples

- ▶ \mathbb{E}^n ;
- ▶ Trees;
- ▶ Non-compact symmetric spaces (e.g. $\mathbb{R}H^2$);
- ▶ Infinite buildings.

Lattices

Let $H = \text{Isom}(X)$ be a locally compact group with Haar measure μ . A discrete subgroup $\Gamma \leq H$ is:

- ▶ a *lattice* if X/Γ has finite covolume;
- ▶ a *uniform lattice* if X/Γ is compact.

For a lattice Γ in a product $\prod_{i=1}^n H_i$ we say Γ is:

- ▶ *irreducible* if the projection to each subproduct of the H_i is non-discrete;
- ▶ *reducible* otherwise.

Examples

- ▶ Crystallographic groups in $\text{Isom}(\mathbb{E}^n)$;
- ▶ Free groups acting on trees in $\text{Aut}(\mathcal{T})$;
- ▶ Arithmetic subgroups of Lie groups e.g. $\text{SL}_2(\mathbb{Z}[\sqrt{2}])$ in $\text{SL}_2(\mathbb{R})^2$;
- ▶ Graph products of finite groups acting on right-angled buildings.

Questions

Let X be a metric space and $H = \text{Isom}(X)$ with Haar measure μ .

1. Does H have lattices?
2. What are properties of a generic lattice in H ?
3. Do the properties of lattices in H reflect properties of H ?
4. Can we classify lattices in H up to isomorphism, commensurability, or isometry?

Trees

A tree \mathcal{T} is a connected graph with no loops.




Figure 1: A 4-regular tree.

Theorem (Bass-Kulkarni 1990)

1. $\text{Aut}(\mathcal{T})$ admits a uniform lattice if and only if \mathcal{T} is the universal cover of a finite connected graph.
2. Lattices in $\text{Aut}(\mathcal{T})$ are fundamental groups of graphs of groups acting faithfully on their Bass-Serre tree. In particular, any uniform lattice is virtually free.

Biautomatic groups

- ▶ An *automatic group* is a finitely generated group equipped with several finite-state automata. These automata represent the Cayley graph of the group. That is, they can tell if a given word representation of a group element is in a "canonical form" and can tell if two elements given in canonical words differ by a generator.
- ▶ A group is *biautomatic* if it has two multiplier automata, for left and right multiplication by elements of the generating set, respectively.

Until 2019 it was not known if every CAT(0) group is biautomatic.

Leary-Minasyan groups

Let $A \in \text{O}(2)$, let L_1 be a finite index subgroup of \mathbb{Z}^2 and let $L_2 = A(L_1)$. Consider the following graph of groups:

$$\mathbb{Z}^2 \curvearrowright L_1^t = L_2$$

We call the fundamental group a *Leary-Minasyan group*. Such a group has the following presentation:

$$\text{LM}(A) = \langle a, b, t \mid [a, b], tL_1t^{-1} = L_2 \rangle$$

The group is equipped with a representation into $\text{Isom}(\mathbb{E}^2)$ which can be described as follows:

$$\phi : \text{LM}(A) \rightarrow \text{Isom}(\mathbb{E}^2) \quad \text{by} \quad \begin{cases} a \mapsto [1, 0]^T \\ b \mapsto [0, 1]^T \\ t \mapsto A. \end{cases}$$

The group acts freely cocompactly on $\mathbb{E}^2 \times \mathcal{T}$ where \mathcal{T} is the Bass-Serre tree. In particular $\text{LM}(A)$ is a CAT(0) group. Note that the construction generalises to $\mathbb{E}^n \times \mathcal{T}$.

Example

Concretely we can take

$$A = \begin{bmatrix} 3/5 & -4/5 \\ 4/5 & 3/5 \end{bmatrix}, \quad L_1 = \left\langle \begin{bmatrix} 2 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\rangle \quad \text{and} \quad L_2 = \left\langle \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix} \right\rangle$$

in this case

$$\text{LM}(A) = \langle a, b, t \mid [a, b], ta^2b^{-1}t^{-1} = a^2b, tab^2t^{-1} = a^{-1}b^2 \rangle.$$




Figure 2: The action of a Leary-Minasyan group on \mathbb{E}^2 . The orthogonal matrix maps the green squares to the blue squares.

Theorem (Leary-Minasyan 2019)

Let $\Gamma = \text{LM}(A)$, \mathcal{T} be the Bass-Serre tree of Γ and $T = \text{Aut}(\mathcal{T})$. Then Γ is virtually biautomatic if and only if A has finite order if and only if Γ is reducible as an $(\text{Isom}(\mathbb{E}^n) \times T)$ -lattice.

Uniform $\text{Isom}(\mathbb{E}^n) \times T$ -lattices

The following lemma gives a rough classification of $\text{Isom}(\mathbb{E}^n) \times T$ -lattices.

Lemma (H. 2021)

Let \mathcal{T} be a locally finite unimodular leafless tree not quasi-isometric to \mathbb{E} and let $T = \text{Aut}(\mathcal{T})$. Every uniform lattice in $\text{Isom}(\mathbb{E}^n) \times T$ splits as a finite graph of virtually abelian groups.

Using the lemma we can prove a number of generic properties for irreducible lattices in $\text{Isom}(\mathbb{E}^n) \times T$.

Theorem (H. 2021)

Let \mathcal{T} be a locally finite unimodular leafless tree not quasi-isometric to \mathbb{E} and let $T = \text{Aut}(\mathcal{T})$. Let Γ be a uniform $(\text{Isom}(\mathbb{E}^n) \times T)$ -lattice. The following are equivalent:

1. Γ is an irreducible $(\text{Isom}(\mathbb{E}^n) \times T)$ -lattice;
2. Γ is irreducible as an abstract group;
3. Γ acts on \mathcal{T} faithfully;
4. Γ does not virtually fibre;
5. Γ is C^* -simple;
6. and if $n = 2$, Γ is non-residually finite and not virtually biautomatic.

The theorem is optimal in the sense that we can show for $n \geq 3$ all irreducible lattices are non-residually finite and not virtually biautomatic. However, there are also reducible lattices with these properties (consider $\mathbb{Z} \times \text{LM}(A)$).

Theorem (H. 2021)

Let $n \geq 2$ and let X be a pentagonal building of thickness $10n$. There exist uniform lattices acting on $\mathbb{E}^n \times X$ which are not virtually biautomatic.

Acknowledgements

I would like to thank my PhD supervisor Professor Ian Leary for his guidance and support.

References

- [1] Hyman Bass and Ravi Kulkarni. Uniform tree lattices. *J. Amer. Math. Soc.*, 3(4):843–902, 1990.
- [2] Hyman Bass and Alexander Lubotzky. *Tree lattices*, volume 176 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 2001. With appendices by Bass, L. Carbone, Lubotzky, G. Rosenberg and J. Tits.
- [3] David B. A. Epstein, James W. Cannon, Derek F. Holt, Silvio V. F. Levy, Michael S. Paterson, and William P. Thurston. *Word processing in groups*. Jones and Bartlett Publishers, Boston, MA, 1992.
- [4] Sam Hughes. Graphs and complexes of lattices. *in preparation*, 2021.
- [5] Ian J. Leary and Ashot Minasyan. Commensurating HNN-extensions: non-positive curvature and biautomaticity. *arXiv:1907.03515 [Math.GR]*, 2019.