

Homological growth of Artin kernels in positive characteristic

Sam P. Fisher¹ · Sam Hughes¹ · Ian J. Leary²

Received: 26 December 2022 / Revised: 26 December 2022 / Accepted: 20 June 2023 / Published online: 6 July 2023 © The Author(s) 2023

Abstract

We prove an analogue of the Lück Approximation Theorem in positive characteristic for certain residually finite rationally soluble (RFRS) groups including right-angled Artin groups and Bestvina–Brady groups. Specifically, we prove that the mod *p* homology growth equals the dimension of the group homology with coefficients in a certain universal division ring and this is independent of the choice of residual chain. For general RFRS groups we obtain an inequality between the invariants. We also consider a number of applications to fibring, amenable category, and minimal volume entropy.

 $\begin{array}{l} \mbox{Mathematics Subject Classification $Primary 20J05 \cdot Secondary $16K99 \cdot 16S35 \cdot 20E26 \cdot 20F36 \cdot 20F65 \cdot 57M07 $} \end{array}$

1 Introduction

A celebrated theorem of Lück relates the rational homology growth in degree *m* through finite covers to the *m*th ℓ^2 -Betti number of a residually finite group. More precisely:

Sam P. Fisher sam.fisher@maths.ox.ac.uk

> Sam Hughes sam.hughes@maths.ox.ac.uk

Ian J. Leary i.j.leary@soton.ac.uk

- ¹ Mathematical Institute, Andrew Wiles Building, Observatory Quarter, University of Oxford, Oxford OX2 6GG, UK
- ² School of Mathematical Sciences, University of Southampton, Southampton SO17 1BJ, UK

Theorem 1.1 (Lück [26]) Let G be a residually finite group of type F_{m+1} and let $(G_n)_{n \in \mathbb{N}}$ be a residual chain of finite index normal subgroups. Then

$$\lim_{n \to \infty} \frac{b_m(G_n; \mathbb{Q})}{[G:G_n]} = b_m^{(2)}(G),$$

where $b_m^{(2)}(G)$ denotes the mth ℓ^2 -Betti number of G.

An immediate consequence of Lück's approximation theorem is that the left-hand limit always exists and is independent of the chosen residual chain. We remind the reader that a *residual chain* is a sequence $G = G_0 \ge G_1 \ge \cdots \ge G_n \ge \cdots$ such that each G_n is a finite index normal subgroup of G and $\bigcap_{n \in \mathbb{N}} G_n = \{1\}$. A related invariant is the *mth* \mathbb{F}_p -homology gradient for a finite field \mathbb{F}_p . It is defined by

$$b_m^{(2)}(G, (G_n); \mathbb{F}_p) := \limsup_{n \to \infty} \frac{b_m(G_n; \mathbb{F}_p)}{[G:G_n]}$$

for any *G* of type F_{m+1} and residual chain $(G_n)_{n \in \mathbb{N}}$. We will recall the definition of various finiteness properties in Sect. 5.

In [27, Conjecture 3.4] Lück conjectured that $b_m^{(2)}(G; \mathbb{F}_p)$ should equal $b_m^{(2)}(G)$ (and hence be independent of the residual chain). This was disproved by Avramidi–Okun–Schreve [1] (using a result of Davis–Leary [13]) where they showed that

$$b_3^{(2)}(A_{\mathbb{R}\mathbf{P}^2}) = 0$$
 but $b_3^{(2)}(A_{\mathbb{R}\mathbf{P}^2}; \mathbb{F}_2) = 1$

independently of the choice of residual chain. Here, $A_{\mathbb{R}\mathbf{P}^2}$ is the right-angled Artin group (RAAG) on (the 1-skeleton of) any flag triangulation of the real projective plane.

For torsion-free groups satisfying the Atiyah conjecture, the ℓ^2 -Betti numbers may be computed via the dimensions of group homology with coefficients in a certain skew field $\mathcal{D}_{\mathbb{Q}G}$, known as the *Linnell skew field* of *G*. In [19], Jaikin-Zapirain introduces analogues of the Linnell skew field with ground ring any skew field \mathbb{F} , denoted $\mathcal{D}_{\mathbb{F}G}$, and called the *Hughes-free division ring* of $\mathbb{F}G$. He proves that $\mathcal{D}_{\mathbb{F}G}$ exists and is unique up to $\mathbb{F}G$ -isomorphism for large classes of groups, including residually finite rationally soluble (RFRS) groups (in particular compact special groups) and conjectures they should exist for all locally indicable groups.

One may compute group homology with coefficients in $\mathcal{D}_{\mathbb{F}G}$ and take $\mathcal{D}_{\mathbb{F}G}$ -dimensions to obtain $\mathcal{D}_{\mathbb{F}G}$ -Betti numbers, denoted $b_m^{\mathcal{D}_{\mathbb{F}G}}(G)$. We emphasise that when G is RFRS and $\mathbb{F} = \mathbb{Q}$, the Linnell skew field and the Hughes free division ring coincide [19, Appendix], and therefore that $b_m^{\mathcal{D}_{\mathbb{Q}G}}(G) = b_m^{(2)}(G)$. The $\mathcal{D}_{\mathbb{F}G}$ -Betti numbers share a number of properties with ℓ^2 -Betti numbers (see for example [18, Theorem 3.9] and [15, Lemmas 6.3 and 6.4]). In light of this we raise the following conjecture:

Conjecture A Let \mathbb{F} be a skew field. Let G be a torsion-free residually finite group of type $\mathsf{FP}_{n+1}(\mathbb{F})$ such that $\mathcal{D}_{\mathbb{F}G}$ exists. Let $(G_i)_{i \in \mathbb{N}}$ be a residual chain of finite index normal subgroups. Then,

$$b_m^{(2)}(G, (G_i); \mathbb{F}) = b_m^{\mathcal{D}_{\mathbb{F}^G}}(G)$$

🖉 Springer

for all $m \leq n$. In particular, the limit supremum in the definition of $b_m^{(2)}$ is a genuine limit and is independent of the choice of residual chain.

Our main result verifies this conjecture for various families of groups—notably for RAAGs, which provided the counterexamples to Lück's original conjecture.

Theorem B Let \mathbb{F} be a skew field and G be a group commensurable with any of

- (1) a residually finite Artin group satisfying the $K(\pi, 1)$ Conjecture, such as a rightangled Artin group or RAAG.
- (2) an Artin kernel, i.e. the kernel of a homomorphism from a RAAG to Z (these include Bestvina–Brady groups);
- (3) a graph product of amenable RFRS groups.

If G is type $\operatorname{FP}_n(\mathbb{F})$, then $b_m^{(2)}(G, (G_n); \mathbb{F}) = b_m^{\mathcal{D}_{\mathbb{F}}G}(G)$ for $m \leq n$. In particular, the limit supremum in the definition of $b_m^{(2)}$ is a genuine limit independent of the choice of residual chain $(G_n)_{n \in \mathbb{N}}$.

In the case when *G* is not torsion-free but contains a finite-index torsion-free subgroup *H*, each of the Betti numbers $b_i(G)$ for *G* appearing in the statement is defined to be $b_i(H)/[G:H]$. This extension to the definition is clearly consistent for $b_i^{(2)}(G; \mathbb{F})$ and is consistent for $b_i^{\mathcal{D}\mathbb{F}G}(G)$ because of [15, Lemma 6.3].

In each case we are able to compute $b_m^{\mathcal{D}_{\mathbb{F}G}}(G)$ explicitly; in cases (1) and (3) we find that it is equal to the \mathbb{F} -homology gradient previously computed in [1] and [30]. In case (2) we compute both the $\mathcal{D}_{\mathbb{F}G}$ -Betti numbers and the \mathbb{F} -homology gradients and show they are equal. We highlight the computation for (2), which we expect will be of independent interest. Note that this generalises the computation of Davis and Okun for the ℓ^2 -Betti numbers of Bestvina–Brady groups [14].

Theorem C Let \mathbb{F} be a skew field, let $\varphi \colon A_L \to \mathbb{Z}$ be an epimorphism and let BB_L^{φ} denote ker φ . If BB_L^{φ} is of type $\mathsf{FP}_n(\mathbb{F})$ then

$$b_m^{\mathcal{D}_{\mathbb{F}^B B_L^{\varphi}}}(BB_L^{\varphi}) = b_m^{(2)}(BB_L^{\varphi}; \mathbb{F}) = \sum_{v \in I^{(0)}} |\varphi(v)| \cdot \widetilde{b}_{m-1}(\mathrm{lk}(v); \mathbb{F}).$$

for all $m \leq n$.

One may extend Conjecture A to G-spaces with finite (n + 1)-skeleton. In this more general setting we are able to verify the conjecture for certain polyhedral product spaces (Theorem 3.15) and certain hyperplane arrangements (Theorem 3.21). We remark that [25] effectively proves the conjecture for torsion-free amenable groups.

For groups where $\mathcal{D}_{\mathbb{F}G}$ exists and is *universal* (see Sect. 4 for a definition), we obtain that the agrarian Betti-numbers give a lower bound for the homology gradients as an easy consequence of a result of Jaikin-Zapirain [19, Corollary 1.6]. Note that the following theorem applies to all residually (amenable and locally indicable) groups by [19, Corollary 1.3], and in particular to RFRS groups.

Theorem D Let \mathbb{F} be a skew-field and let G be a residually finite group of type $\mathsf{FP}_{n+1}(\mathbb{F})$ such that $\mathcal{D}_{\mathbb{F}G}$ exists and is the universal division ring of fractions of $\mathbb{F}G$. Then

$$b_m^{\mathcal{D}_{\mathbb{F}^G}}(G) \le b_m^{(2)}(G, (G_i); \mathbb{F})$$

for all $m \leq n$, where $(G_i)_{i \in \mathbb{N}}$ is any residual chain of finite-index subgroups of G.

We also mention the work of Bergeron–Linnell–Lück–Sauer [4], which we believe provides some more evidence for Conjecture A. Let Γ be the fundamental group of a finite CW complex X with a homomorphism $\varphi \colon \Gamma \to \operatorname{GL}_n(\mathbb{Z}_p)$. Let G be the closure of $\varphi(\Gamma)$ and let $G_i = \operatorname{ker}(G \to \operatorname{GL}_n(\mathbb{Z}/p^i\mathbb{Z}))$. Recall that the *Iwasawa algebra* of G over \mathbb{F}_p is $\mathbb{F}_p[\![G]\!] = \varprojlim_p \mathbb{F}_p[G/G_i]$. If G is torsion-free, then $\mathbb{F}_p[\![G]\!]$ has no zero divisors and is Ore with respect to its nonzero elements. Letting $\overline{\Gamma} = \Gamma/\operatorname{ker}\varphi$, we have ring homomorphisms

$$\mathbb{F}\overline{\Gamma} \to \mathbb{F}G \to \mathbb{F}_p[\![G]\!] \to \mathcal{D},$$

where \mathcal{D} is the Ore localisation of $\mathbb{F}_p[\![G]\!]$. If M is an $\mathbb{F}_p[\![G]\!]$ -module and G is torsion-free, then the dimension of M is

$$\dim_{\mathbb{F}_p\llbracket G\rrbracket} M := \dim_{\mathcal{D}}(\mathcal{D} \otimes_{\mathbb{F}_p\llbracket G\rrbracket} M).$$

If *G* is not torsion-free, then we can pass to a uniform finite-index subgroup $G_0 \le G$ and then define dim_{**F**_p[G]} $M = [G : G_0]^{-1} \dim_{$ **F** $_p[G]} M$. There is then a natural mod *p* analogue of ℓ^2 -Betti numbers given by

$$\beta_k(\overline{X},\overline{\Gamma};\mathbb{F}_p) = \dim_{\mathbb{F}_p[\![G]\!]} H_k(\mathbb{F}_p[\![G]\!] \otimes_{\mathbb{F}_p\overline{\Gamma}} C_{\bullet}(\overline{X};\mathbb{F}_p))$$

where \overline{X} is the cover of X corresponding to ker φ . With this setup, Bergeron–Linnell–Lück–Sauer prove the following mod-*p* Lück approximation style theorem.

Theorem 1.2 With the notation above, let $\Gamma_i = \varphi^{-1}(G_i)$ and let X_i be the corresponding cover of X. Then

$$b_k(X_i; \mathbb{F}_p) = [\Gamma : \Gamma_i] \cdot \beta_k(\overline{X}, \overline{\Gamma}; \mathbb{F}_p) + O\left([\Gamma : \Gamma_i]^{1 - \frac{1}{\dim G}}\right)$$

for all k.

1.1 Outline of the paper

In Sect. 2 we give the relevant background on group rings and the computational tools we will need. In fact in many computations we are able to work in the more general setting of *agrarian invariants* as defined in [18] and so summarise the relevant theory. The remaining three sections of the paper are described in the sequel.

In Sect. 3 we introduce the notion of a *confident complex*; roughly this is a CW complex admitting a finite cover by subcomplexes with amenable fundamental group such that the nerve of the cover has good properties. We then compute the $\mathcal{D}_{\mathbb{F}G}$ -Betti numbers and \mathbb{F}_p -homology gradients of these complexes showing they are related to \mathbb{F}_p -Betti numbers of the nerve. The computation of the first invariant a uses a spectral sequence collapsing result of Davis–Okun [14] building on work of Davis–Leary [13]. The computation of the second invariant is similar in spirit to the work of Avramidi–Okun–Schreve [1] and Okun–Schreve [30] and again relies on a spectral sequence argument.

The remainder of the section involves computations of the homological invariants for various spaces and groups with the goal of showing they satisfy Conjecture A. The computations are summarised as follows. In Theorem 3.12 we show that Artin kernels of type FP_n are fundamental groups of confident spaces with an *n*-acyclic covering space, and use this to prove Theorem C. In Theorem 3.15 we compute the invariants for graph products of amenable RFRS groups (including RAAGs) and polyhedral products of classifying spaces of amenable RFRS groups. In Theorem 3.19 we compute the invariants for the Artin groups alluded to earlier. In fact the method applies to RFRS groups admitting a strict fundamental domain with certain stabilisers (Remark 3.20). Finally, inspired by [12], we compute the invariants for hyperplane arrangements whenever the invariants are defined (Theorem 3.21).

1.1.2 A lower bound for homology gradients

In Sect. 4, we recall the notion of a universal division ring of fractions and prove Theorem D as a consequence of work of Jaikin-Zapirain in [19].

1.1.3 Applications to fibring

In Theorem 5.1 we prove that if a RFRS group of type $\mathsf{FP}_{n+1}(\mathbb{F})$ is not virtually $\mathsf{FP}_n(\mathbb{F})$ -fibred, then there is some $m \leq n$ such that $b_m^{(2)}(G, (G_n); \mathbb{F}) > 0$ for every residual chain $(G_n)_{n \in \mathbb{N}}$.

In the remainder of Sect. 5 we apply the computations of the agrarian invariants of RAAGs and Artin kernels to obtain some results about fibring in RAAGs. In particular, we make progress towards the following question of Matthew Zaremsky, communicated to us by Robert Kropholler: If a RAAG virtually algebraically fibres with kernel of type F_n , then does it algebraically fibre with kernel of type F_n ? We are able to answer this question if one replaces F_n with $FP_n(R)$, where R is either a skew field, \mathbb{Z} , or \mathbb{Z}/m for some integer m > 1 (Theorem 5.2). Since finitely presented groups of type FP_n are of type F_n , this leaves F_2 as the main case of interest in Zaremsky's question.

In [15], the first author showed that if *G* is RFRS and ℓ^2 -acyclic in dimensions $\leq n$, then *G* virtually $\mathsf{FP}_n(\mathbb{Q})$ -fibres, but left unanswered whether ℓ^2 -Betti numbers control virtual FP_n -fibring. We resolve that here by showing that there are RAAGs that are ℓ^2 -acyclic but do not virtually FP_2 -fibre (Proposition 5.5). Finally, we use the

explicit computation of the agrarian invariants of Artin kernels to show that if A_L is a RAAG and \mathbb{F} is a skew field, then either all of the FP_n(\mathbb{F})-fibres of A_L are themselves virtually FP_n(\mathbb{F})-fibred or none of them are (Theorem 5.6, Corollary 5.8).

1.1.4 Applications to amenable category and minimal volume entropy

In Sect. 6 we relate the Agrarian invariants to the amenable category of [9] and the minimal volume entropy of Gromov [16]. The relevant background is described in the section.

We show via an argument of Sauer [31] that having a small amenable category implies vanishing of $\mathcal{D}_{\mathbb{F}G}$ -Betti numbers for residually finite groups (Proposition 6.1). We also show that for residually finite groups with uniformly uniform exponential growth admitting a finite K(G, 1), having a non-zero $\mathcal{D}_{\mathbb{F}G}$ -Betti number implies the minimal volume entropy of *G* is non-zero (Corollary 6.2). The analogous results for \mathbb{F}_p -homology gradients were established by Sauer [31] and Haulmark–Schreve [17] respectively. In some sense this provides more evidence towards Conjecture A. Finally, we give a condition for the minimal volume entropy of an Artin kernel admitting a finite K(G, 1) to be non-zero (Corollary 6.3) and conjecture a converse.

2 Preliminaries

Throughout all rings are assumed to be associative and unital.

2.1 Finiteness properties

Let *R* be a ring and *G* be a group. Then *G* is said to be of *type* $\text{FP}_n(R)$ if there is a projective resolution $P_{\bullet} \to R$ of the trivial *RG*-module *R* such that P_i is finitely generated for all $i \leq n$. If *G* is of type $\text{FP}_n(R)$ and *S* is an *R*-algebra, then *G* is of type $\text{FP}_n(S)$. Thus, if *G* is of type $\text{FP}_n(\mathbb{Z})$, then *G* is of type $\text{FP}_n(R)$ for any ring *R*; because of this, we write FP_n to mean $\text{FP}_n(\mathbb{Z})$. Note that finite generation is equivalent to $\text{FP}_1(R)$ for any ring *R*, though FP_2 is in general a stronger condition than $\text{FP}_2(R)$.

We also the mention the homotopical analogue of the $\mathsf{FP}_n(R)$ condition: a group G is of type F_n if G has a classifying space with finite *n*-skeleton. Note that F_1 is equivalent to $\mathsf{FP}_1(R)$ for any ring R, but that F_n is in general strictly stronger than $\mathsf{FP}_n(R)$ for $n \ge 2$.

2.2 Agrarian invariants

Let *G* be a group and let \mathbb{F} be a skew field. The *group ring* $\mathbb{F}G$ is the set of formal sums $\sum_{g \in G} \lambda_g g$, where $\lambda_g \in \mathbb{F}$ is zero for all but finitely many $g \in G$, equipped with the obvious addition and multiplication operations. Let \mathcal{D} be a skew field. Then an *agrarian embedding* is a ring monomorphism $\alpha : \mathbb{F}G \hookrightarrow \mathcal{D}$. Agrarian embeddings were first studied by [28, 29], who proved that group rings of bi-orderable groups have agrarian embeddings. Note that the existence of an agrarian embedding implies that

G is torsion-free and that the Kaplansky zero divisor and idempotent conjectures hold for $\mathbb{F}G$. There is no known example of a torsion-free group *G* and a skew field \mathbb{F} such that $\mathbb{F}G$ does not have an agrarian embedding.

Let *X* be a CW-complex with a cellular *G* action such that for every $g \in G$ and every open cell *e* of *X*, if $g \cdot e \cap e \neq \emptyset$ then *g* fixes *e* pointwise. The cellular chain complex $C_{\bullet}(X; \mathbb{F})$ is naturally an $\mathbb{F}G$ -module. In this situation, *X* is called a *G*-CW complex. If $\alpha : \mathbb{F}G \to \mathcal{D}$ is an agrarian map, then we can define the \mathcal{D} -homology and \mathcal{D} -Betti numbers of *X* by

$$H_p^{\mathcal{D}}(X) := H_p(\mathcal{D} \otimes_{\mathbb{F}G} C_{\bullet}(X))$$
 and $b_p^{\mathcal{D}}(X) = \dim_{\mathcal{D}} H_p^{\mathcal{D}}(X)$

where dim $_{\mathcal{D}}$ denotes the dimension as a \mathcal{D} -module, which is well-defined since \mathcal{D} is a skew field. Taking X to be a classifying space of G, we obtain the \mathcal{D} -homology and \mathcal{D} -Betti numbers of G.

The following theorem gives a central example of an agrarian embedding.

Theorem 2.1 (Linnell [24]) *If G is a torsion-free group satisfying the* strong Atiyah conjecture, *then there is a skew field* $\mathcal{D}_{\mathbb{Q}G}$, *known as the* Linnell skew field *of G*, *and an agrarian embedding* $\mathbb{Q}G \hookrightarrow \mathcal{D}_{\mathbb{Q}G}$ *such that* $b_p^{(2)}(G) = b^{\mathcal{D}_{\mathbb{Q}G}}(G)$.

The strong Atiyah conjecture asserts that if *X* is a free *G*-CW complex of finite type and *G* has finite subgroups of bounded order, then $lcm(G) \cdot b_p^{(2)}(X) \in \mathbb{Z}$ for all $p \in \mathbb{N}$, where lcm(G) is the least common multiple of the orders of finite subgroups of *G*. The strong Atiyah conjecture is known for many groups, in particular for residually (torsion-free solvable groups) [32], for cocompact special groups [33], and for locally indicable groups [20]. Importantly for us, the Atiyah conjecture holds for all RFRS groups, and in particular all subgroups of RAAGs. The following theorem of Jaikin-Zapirain provides many examples of agrarian embeddings in positive characteristic.

Theorem 2.2 (Jaikin-Zapirain [19, Theorem 1.1]) Let \mathbb{F} be a skew field and let G be either locally indicable amenable, residually (torsion-free nilpotent), or free-bycyclic. Then there exists a division ring $\mathcal{D}_{\mathbb{F}G}$, known as the Hughes-free division ring of $\mathbb{F}G$, and an agrarian embedding $\mathbb{F}G \hookrightarrow \mathcal{D}_{\mathbb{F}G}$.

2.3 A Mayer–Vietoris type spectral sequence

The following construction of a Mayer–Vietoris type spectral sequence is due to Davis and Okun [14]. We will state the homological version of the spectral sequence with arbitrary coefficients. Let P be a poset. We define Flag(P) to be the simplicial realisation of P, i.e. Flag(P) is the simplicial complex whose simplices are the totally ordered, finite, nonempty subsets of P. Hence, every simplex $\sigma \in$ Flag(P) has a well-defined *minimum vertex*, denoted min(σ).

If Y is a CW complex, then a *poset of spaces* in Y over P is a cover $\mathcal{Y} = \{Y_a\}_{a \in P}$ of Y with each Y_a a subcomplex such that

(1) a < b implies $Y_a \subseteq Y_b$;

(2) \mathcal{Y} is closed under finite, nonempty intersections.

Let *R* be a ring and let Mod_R denote the category *R*-modules. A *poset of coefficients* for P is a contravariant functor $\mathcal{A}: P \to Mod_R$. The functor \mathcal{A} induces a system of coefficients on Flag(P) via $\sigma \mapsto \mathcal{A}_{\min(\sigma)}$, which gives chain complex

$$C_j(\operatorname{Flag}(\mathsf{P}); \mathcal{A}) := \bigoplus_{\sigma \in \operatorname{Flag}(\mathsf{P})^{(j)}} \mathcal{A}_{\min(\sigma)},$$

where $\operatorname{Flag}(\mathsf{P})^{(j)}$ is the set of *j*-simplices in $\operatorname{Flag}(\mathsf{P})$.

Lemma 2.3 ([14, Lemmas 2.1 and 2.2]) Let M be an R-module and suppose $\mathcal{Y} = \{Y_a\}_{a \in \mathsf{P}}$ is a poset of spaces over P in a CW complex Y. There is a Mayer–Vietoris type spectral sequence

$$E_{p,q}^2 = H_p(\operatorname{Flag}(\mathsf{P}); \mathcal{H}_q(\mathcal{Y}; M)) \to H_{p+q}(Y; M),$$

where $\mathcal{H}_{\bullet}(\mathcal{Y}; M)$ is a system of coefficients given by

$$\mathcal{H}_{\bullet}(\mathcal{Y}; M)(\sigma) = H_{\bullet}(Y_{\min(\sigma)}; M)$$

Moreover, if the induced homomorphism $H_{\bullet}(Y_a; M) \to H_{\bullet}(Y_b; M)$ is zero whenever a < b in P, then

$$E_{p,q}^{2} = \bigoplus_{a \in \mathsf{P}} H_{p}(\operatorname{Flag}(\mathsf{P}_{\geq a}), \operatorname{Flag}(\mathsf{P}_{>a}); H_{q}(Y_{a}; M)).$$

3 Computations

3.1 Approximation for spaces with confident covers

Fix a skew field \mathbb{F} and let *X* be a compact CW complex with a finite poset of spaces $\mathcal{X} = \{X_{\alpha}\}_{\alpha \in \mathsf{P}}$ over P . We call \mathcal{X} confident if it satisfies the following conditions: for each $\alpha \in \mathsf{P}$

- (i) each X_{α} has finitely many components;
- (ii) either $X_{\alpha} \subseteq X^{(0)}$ or each connected component of X_{α} is a classifying space with torsion-free amenable fundamental group such that $\mathbb{F}G$ has no zero-divisors;
- (iii) if $C \subseteq X_{\alpha}$ is a component, then the inclusion $C \subseteq X$ induces an injection $\pi_1(C) \to \pi_1(X)$;
- (iv) if X_{α} is a collection of points, then α is minimal in P; equivalently, $X_{\alpha} \cap X_{\beta} = \emptyset$ whenever $X_{\alpha}, X_{\beta} \subseteq X^{(0)}$.

Remark 3.1 If G is a torsion-free elementary amenable group, then $\mathbb{F}G$ has no zerodivisors.

🖄 Springer

We will show that spaces with confident covers satisfy Lück's approximation theorem in arbitrary characteristic, following Avramidi, Okun, and Schreve who prove the same result for the Salvetti complex of a RAAG. We will then show that the result agrees with the agrarian Betti numbers of the space.

Before beginning, we fix some notation. For the rest of the section, *X* will be a compact CW complex with a confident cover $\mathcal{X} = \{X_{\alpha}\}_{\alpha \in \mathsf{P}}$ and we assume that $G := \pi_1(X)$ is residually finite with residual chain $(G_n)_{n \in \mathbb{N}}$. Let X_n be the covering space of *X* corresponding to $G_n \leq G$. Fix some $n \in \mathbb{N}$ and let $V = \mathbb{F}[G/G_n]$. Let *K* be the nerve of \mathcal{X} and let $X_{\sigma} = X_{\min(\sigma)}$ for any simplex $\sigma \in K$. By an abuse of notation, we will use α to denote both an element of P and the corresponding vertex of *K*.

Recall the Mayer-Vietoris type spectral sequence

$$E_{p,q}^{1} = C_{p}(K; H_{q}(X_{\sigma}; V)) \Rightarrow H_{p+q}(X; V) \cong H_{p+q}(X_{n}; \mathbb{F})$$

(see, e.g., [7, VII.4]).

Lemma 3.2 We have that

$$\lim_{n \to \infty} \frac{\dim_{\mathbb{F}} H_q(X_{\sigma}; V)}{[G:G_n]} = \begin{cases} n_{\sigma} & \text{if } q = 0 \text{ and } X_{\sigma} \subseteq X^{(0)}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof The proof is similar to that of [1, Lemma 8]. The claim is clear when X_{σ} consists of 0-cells. In the other case, since the homology growth of amenable groups satisfying (ii) is sublinear [25, Theorem 0.2], the only way dim_{\mathbb{F}} $H_q(X_{\sigma}; V)$ can grow linearly is if the number of components of the preimage of X_{σ} in X_n grows linearly with the index. But this does not occur since the sequence Γ_n is residual and normal and the inclusions $X_{\sigma} \subseteq X$ induce π_1 -injections of infinite groups on each component of X_{α} .

The spectral sequence is therefore concentrated on the $E_{p,0}^1$ line, up to an error sublinear in the index [$G : G_n$]. This implies that

$$\limsup_{n \in \mathbb{N}} \frac{\dim_{\mathbb{F}} E_{p,0}^2}{[G:G_n]} = \limsup_{n \in \mathbb{N}} \frac{\dim_{\mathbb{F}} H_p(X;V)}{[G:G_n]}.$$
 (1)

Define a poset of coefficients on the vertices of K by

$$\mathcal{A}_{\alpha} = \begin{cases} V^{n_{\sigma}} & \text{if } X_{\sigma} \subseteq X^{(0)}, \\ 0 & \text{otherwise.} \end{cases}$$

Let $K^{(p)}$ denote the set of *p*-simplices of *K*. There is a chain projection

$$E_{p,0}^{1} = \bigoplus_{\sigma \in K^{(p)}} V^{n_{\sigma}} \to C_{p}(K; \mathcal{A}_{\alpha}) = \bigoplus_{\sigma \in K^{(p)}: X_{\min(\sigma)} \subseteq X^{(0)}} V^{n_{\sigma}}.$$

By Lemma 3.2, the kernel of this projection has dimension sublinear in the index $[G:G_n]$ and therefore

$$\limsup_{n \in \mathbb{N}} \frac{\dim_{\mathbb{F}} E_{p,0}^2}{[G:G_n]} = \limsup_{n \in \mathbb{N}} \frac{\dim_{\mathbb{F}} H_p(X; \mathcal{A}_{\alpha})}{[G:G_n]}.$$
 (2)

The proof of the following proposition is similar to that of [1, Lemma 9], except that in their case the nerve is contractible. Though *K* is not necessarily contractible, it does decompose nicely into contractible pieces centred at the vertices $\alpha \in K^{(0)}$ such that $X_{\alpha} \subseteq X^{(0)}$.

Proposition 3.3 Let $S = \{ \alpha \in K^{(0)} : X_{\alpha} \subseteq X^{(0)} \}$. Then

$$\dim_{\mathbb{F}} H_p(K; \mathcal{A}_{\sigma}) = [G: G_n] \cdot \sum_{\alpha \in S} n_{\alpha} \widetilde{b}_{p-1}(\mathrm{lk}(\alpha); \mathbb{F})$$

In particular, $\lim_{n\to\infty} \frac{b_p(X_n;\mathbb{F})}{[G:G_n]}$ exists and is independent of the residual chain (G_n) .

Proof By (iv), if $\alpha \in K^{(0)}$ is a vertex such that $\mathcal{A}_{\alpha} \neq 0$, then every vertex $\beta \in K^{(0)}$ adjacent to α has $\mathcal{A}_{\beta} = 0$. Therefore the chain complex $C_{\bullet}(K; \mathcal{A}_{\sigma})$ decomposes as a direct sum of chain complexes $\bigoplus_{\alpha \in S} C_{\bullet}(\operatorname{st}(\alpha); \mathcal{A}_{\sigma})$, where the coefficient system \mathcal{A}_{σ} is restricted to each st $(\alpha) \subseteq K$.

For each $\alpha \in S$, there is a short exact sequence of chain complexes

$$0 \to C_{\bullet}(\mathrm{lk}(\alpha); \mathbb{F}) \otimes V^{n_{\alpha}} \to C_{\bullet}(\mathrm{st}(\alpha); \mathbb{F}) \otimes V^{n_{\alpha}} \to C_{\bullet}(\mathrm{st}(\alpha); \mathcal{A}_{\sigma}) \to 0.$$

Because $st(\alpha)$ is contractible, the middle term is acyclic and therefore

$$H_{\bullet}(\mathrm{st}(\alpha); \mathcal{A}_{\sigma}) \cong H_{\bullet-1}(\mathrm{lk}(\alpha); \mathbb{F}) \otimes V^{n_{\alpha}}.$$

The formula in the statement of the proposition follows, since $V^{n_{\alpha}}$ is a vector space of dimension $[G:G_n] \cdot n_{\alpha}$.

This formula together with (1) and (2) show that $\limsup_n \frac{b_p(X_n;\mathbb{F})}{[G:G_n]}$ is independent of the residual chain (G_n) . Moreover, this implies that the lim sup is a genuine limit.

Remark 3.4 In [1], the computation of Proposition 3.3 is carried out in the case that X_L is the Salvetti complex of the RAAG determined by L. If \mathcal{X} is the cover of X_L by standard tori, then there is a single vertex v in the cover and the corresponding vertex α in the nerve has link isomorphic to L. Thus, we recover the formula $b_p^{(2)}(A_L; \mathbb{F}) = \tilde{b}_{p-1}(L; \mathbb{F})$.

Remark 3.5 Condition (ii) can be weakened as follows: One only requires that first, $\pi_1 X$ is residually finite; and second, that each X_{σ} is (homotopy equivalent to) a compact CW complex with vanishing \mathbb{F} - ℓ^2 -Betti numbers independent of the chain, or is a 0-cell. In this case the conclusion of Proposition 3.3 still holds.

Corollary 3.6 Let X be a confident CW complex with $\pi_1(X)$ residually finite. If $\pi: \widehat{X} \to X$ is a degree d cover, then $b_p^{(2)}(H; \mathbb{F}) = d \cdot b_p^{(2)}(G; \mathbb{F})$.

Proof Let $\{X_{\alpha}\}$ be a confident cover of X. Then $\{\pi^{-1}(X_{\alpha})\}$ is a confident cover of \widehat{X} and $|\pi^{-1}(X_{\alpha})| = d \cdot |X_{\alpha}|$ whenever $X_{\alpha} \subseteq X^{(0)}$.

3.2 Agrarian homology of spaces with confident covers

We continue with the same set-up as the previous subsection: *X* is a CW complex with a confident cover $\mathcal{X} = \{X_{\alpha}\}_{\alpha \in \mathsf{P}}$ and let *K* be the covering. Notice that $K \cong \mathrm{Flag}(\mathsf{P})$. Additionally, we will assume that there exists a skew field \mathcal{D} and a fixed agrarian embedding $\mathbb{F}G \to \mathcal{D}$, where $G = \pi_1(X)$.

Proposition 3.7 *Let* $S = \{ \alpha \in K^{(0)} : X_{\alpha} \subseteq X^{(0)} \}$ *. Then*

$$\dim_{\mathbb{F}} H_p^{\mathcal{D}}(X) = \sum_{\alpha \in S} n_{\alpha} \widetilde{b}_{p-1}(\mathrm{lk}(\alpha); \mathbb{F}).$$

In particular, if $\pi_1(X)$ is residually finite then $b_p^{\mathcal{D}}(X) = b_p^{(2)}(X; \mathbb{F})$.

Proof Suppose X_{α} does not consist of 0-cells. Then, each component of X_{α} is a classifying space for an infinite amenable group and therefore $H_p^{\mathcal{D}}(X_{\alpha}) = 0$ by [18, Theorem 3.9(6)]. Since the elements α such that $X_{\alpha} \subseteq X^{(0)}$ are minimal in P, the spectral sequence of Lemma 2.3 collapses on the $E_{p,0}^2$ line. By Lemma 2.3,

$$H_p^{\mathcal{D}}(X) = \bigoplus_{\alpha \in S} H_p(\operatorname{Flag}(\mathsf{P}_{\geq \alpha}), \operatorname{Flag}(\mathsf{P}_{>\alpha}); \mathcal{D}^{n_\alpha}),$$

whence the stated formula follows.

Corollary 3.8 If X is confident, $G = \pi_1(X)$ is residually finite, and G has a Hughesfree division ring $\mathcal{D}_{\mathbb{F}G}$, then $b_p^{\mathcal{D}_{\mathbb{F}H}} = b_p^{(2)}(\widehat{X}; \mathbb{F})$ for every finite index subgroup $H \leq G$ and corresponding finite covering space $\widehat{X} \to X$.

Proof This follows from Corollary 3.6 and the fact that Hughes-free Betti numbers scale when passing to a finite index subgroup [15, Lemma 6.3].

Remark 3.9 Condition (ii) can be weakened as follows: One only requires that first $\pi_1 X$ admits an agrarian map $\pi_1 X \to \mathcal{D}$; and second that each X_{σ} is (homotopy equivalent to) a compact CW complex with with vanishing \mathcal{D} -Betti numbers or is a 0-cell. In this case the conclusion of Proposition 3.7 still holds.

3.3 Artin kernels

An Artin kernel is simply the kernel of a non-zero homomorphism $\varphi \colon A_L \to \mathbb{Z}$, where *L* is a flag complex and A_L is the RAAG it determines. We now apply the results of

the previous two subsections to Artin kernels and obtain an explicit formula for their agrarian Betti numbers in terms of L and φ .

We fix a skew field \mathbb{F} , a flag complex L, and a surjective homomorphism $\varphi \colon A_L \to \mathbb{Z}$, where A_L is the RAAG on L. Moreover, we fix a standard generating set for A_L , identified with the vertex set Vert(L). We denote the Artin kernel by $BB_L^{\varphi} := \ker \varphi$. If φ is the map sending each of the generators to $1 \in \mathbb{Z}$, then $BB_L^{\varphi} = BB_L$ is the usual Bestvina–Brady group.

Let T_L be the Salvetti complex on L and let X_L be its universal cover. There is an affine map $T_L \to S^1$ inducing φ constructed as follows. For each $v \in \text{Vert}(L)$, let $S_v^1 := \mathbb{R}/\mathbb{Z}$ be the corresponding circle in T_L . Let $\sigma = \{v_1, \ldots, v_k\} \in L$ be a simplex and let $T_\sigma = S_{v_1}^1 \times \cdots \times S_{v_k}^1$ be the associated subtorus of T_L . There is a map

 $T_{\sigma} \to S^1 = \mathbb{R}/\mathbb{Z}, \quad (x_1, \dots, x_k) \longmapsto \varphi(v_1)x_1 + \dots + \varphi(v_k)x_k + \mathbb{Z}.$

The maps on each of the subtori extend to a well-defined map $f: T_L \to S^1$ inducing φ on the level of fundamental groups. Moreover, f induces a cube-wise affine A_L -equivariant height function $h: X_L \to \mathbb{R}$ making the diagram

$$\begin{array}{ccc} X_L & \stackrel{h}{\longrightarrow} & \mathbb{R} \\ \downarrow & & \downarrow \\ T_L & \stackrel{f}{\longrightarrow} & S^1 \end{array}$$

commute—here the vertical arrows are the universal covering maps.

We borrow the following definition and terminology from [8].

Definition 3.10 A vertex v of L is *living* [resp., *dead*] if $\varphi(v) \neq 0$ [resp., $\varphi(v) = 0$]. Denote the full subcomplex of L spanned by the living [resp., dead] vertices by L^a [resp., L^d].

Let $Z = h^{-1}(\{p\})$ for some $p \notin \mathbb{Z}$. The level set Z has a natural CW complex structure and BB_L^{φ} acts cocompactly on Z; we denote the quotient Z/BB_L^{φ} by Y. For each *n*-simplex $\sigma \in L$, the subtorus $T_{\sigma} \subseteq T_L$ lifts to X_{σ} , a collection of pairwise disjoint *sheets* in X_L . Each sheet is an isometrically embedded copy of (n + 1)-dimensional Euclidean space.

Let P be the poset of simplices of L that contain at least one vertex in L^a . Then Z is covered by the collection $\{X_{\sigma} \cap Z\}_{\sigma \in P}$. Writing Y_{σ} for the image of $X_{\sigma} \cap Z$ in Y, we obtain a poset of spaces $\mathcal{Y} = \{Y_{\sigma}\}_{\sigma \in P}$ of Y where each subcomplex Y_{σ} is a disjoint union of tori or a set of vertices. Crucially, \mathcal{Y} is a confident cover.

Lemma 3.11 If $\sigma = \{v\} \in L^a$ is a vertex, then Y_{σ} is a collection of $|\varphi(v)|$ vertices.

Proof We will show that there are exactly $|\varphi(v)|$ orbits of lines in X_{σ} under the BB_{L}^{ρ} -action on X_{L} . For each vertex v in L^{a} , evenly subdivide each edge of X_{v} into $|\varphi(v)|$ segments. Note that the restriction $\overline{X}_{L}^{(1)} \rightarrow \mathbb{R}$ of h is cellular, where $\overline{X}_{L}^{(1)}$ is the subdivided 1-skeleton of X_{L} and \mathbb{R} is given the cell structure with vertex set \mathbb{Z} and edge set $\{[n, n + 1] : n \in \mathbb{Z}\}$.

Fix a vertex $\sigma = \{v\} \in L^a$ and let \overline{X}_{σ} be the subdivision of X_{σ} . Let *e* be an edge of \overline{X}_{σ} and let *e'* be the unique edge of X_{σ} such that $e \subseteq e'$. We say *e* is an edge of *type i* if it is the *i*th highest edge (under the height function *h*) contained in *e'*; the integer *i* can take values in $\{0, \ldots, |\varphi(v)| - 1\}$.

Because the action of BB_L^{φ} on X_L is height preserving, it preserves the set of type i edges. Since φ is surjective, $gcd((\varphi(v))_{v \in L^{(0)}}) = 1$. Therefore, the generic level set Z intersects edges of type i for every $i \in \{0, \ldots, |\varphi(v)| - 1\}$. Moreover, BB_L^{φ} acts transitively on the set of edges of type i of the same height, which follows from the fact that BB_L^{φ} acts transitively on the set of edges of T_{σ} of the same height. Thus, we conclude that there are exactly $|\varphi(v)|$ orbits of lines in X_{σ} under the BB_L^{φ} action. \Box

Theorem 3.12 Let $\varphi \colon A_L \to \mathbb{Z}$ be an epimorphism and let Y be a generic level set of the induced height function. If $\mathbb{F}BB_L^{\varphi} \to \mathcal{D}$ is an agrarian embedding, then

$$b_p^{\mathcal{D}}(Y) = b_p^{(2)}(Y; \mathbb{F}) = \sum_{v \in L^{(0)}} |\varphi(v)| \cdot \widetilde{b}_{p-1}(\operatorname{lk}(v); \mathbb{F}).$$

Moreover, if BB_I^{φ} is of type $FP_n(\mathbb{F})$ then

$$b_p^{\mathcal{D}}(BB_L^{\varphi}) = b_p^{(2)}(BB_L^{\varphi}; \mathbb{F}) = \sum_{v \in L^{(0)}} |\varphi(v)| \cdot \widetilde{b}_{p-1}(\operatorname{lk}(v); \mathbb{F})$$

for all $p \leq n$. We also have $b_p^{\mathcal{D}}(H) = b_p^{(2)}(H; \mathbb{F})$ whenever H is a finite index subgroup of BB_I^{φ} and \mathcal{D} is the Hughes-free division ring of $\mathbb{F}H$.

Before beginning the proof, we remark that BB_L^{φ} is a RFRS group, being a subgroup of a RAAG. Hence, by [19, Corollary 1.3], $\mathcal{D}_{\mathbb{F}BB_L^{\varphi}}$ exists. When $\mathbb{F} = \mathbb{Q}$, the Hughes free division ring and the Linnell skew field coincide, so in this case Theorem 3.12 computes the ℓ^2 -Betti numbers of BB_L^{φ} . This generalises the computation of Davis and Okun in [14, Theorem 4.4]

Proof The first statement is an immediate consequence of Propositions 3.3 and 3.7, Remark 3.9, Remark 3.5, Lemma 3.11, and the observation that \mathcal{Y} is a confident cover. The second follows from the fact that if BB_L^{φ} is of type $FP_n(\mathbb{F})$ if and only if Z is *n*-acyclic with \mathbb{F} coefficients [8] (Bux–Gonzalez consider only the case $\mathbb{F} = \mathbb{Z}$, but their result remains true over arbitrary coefficients).

Remark 3.13 The condition that BB_L^{φ} is of type $\mathsf{FP}_n(\mathbb{F})$ can be verified directly in the flag complex *L*. Recall that a topological space is *n*-acyclic if its reduced homology (with coefficients in \mathbb{F} in our case) vanishes in degrees $\leq n$. Note that we use the convention that the reduced homology of the empty set is \mathbb{F} in dimension -1, so if *X* is *n*-acyclic for $n \geq -1$, then *X* is nonempty.

Bux and Gonzalez show that BB_L^{φ} is of type $\mathsf{FP}_n(\mathbb{F})$ if and only if $L^a \cap \mathrm{lk}(\sigma)$ is $(n - \dim(\sigma) - 1)$ -acyclic (with coefficients in \mathbb{F}) for every simplex $\sigma \in L^d$, including the empty simplex which has dimension -1 and link L [8, Theorem 14]. Their result is stated in the case $\mathbb{F} = \mathbb{Z}$ but remains true, with the same proof, when stated over a

general coefficient ring. We will use this characterisation of the finiteness properties of BB_I^{φ} is Sect. 5.

It is known by work of Okun–Schreve [30] that for RAAGs the homology torsion growth $t_p^{(2)}(A_L)$ in degree p is equal to $|H_{p-1}(L; \mathbb{Z})_{tors}|$. We conjecture an analogous result for Artin kernels.

Conjecture 3.14 If BB_L^{φ} is of type FP_n , then

$$t_p^{(2)}(BB_L^{\varphi}) = \limsup_{n \to \infty} \frac{\log |H_p(G_n; \mathbb{Z})_{\text{tors}}|}{[BB_L^{\varphi}:G_n]} = \sum_{v \in L^{(0)}} |\varphi(v)| \cdot |H_{p-1}(\operatorname{lk}(v); \mathbb{Z})_{\text{tors}}|$$

for $p \leq n$ and any residual chain (G_n) .

3.4 Graph products

Let *K* be a simplicial complex on the vertex set $[m] := \{1, ..., m\}$. Let $(\mathbf{X}, \mathbf{A}) = \{(X_i, A_i) : i \in [m]\}$ be a collection of CW-pairs. The *polyhedral product* of (\mathbf{X}, \mathbf{A}) and *K* is the space

$$(\mathbf{X}, \mathbf{A})^K := \bigcup_{\sigma \in K} \prod_{i=1}^m Y_i^{\sigma} \text{ where } Y_i^{\sigma} = \begin{cases} X_i & \text{if } i \in \sigma, \\ A_i & \text{if } i \notin \sigma. \end{cases}$$

If $\Gamma = {\Gamma_1, ..., \Gamma_m}$ is a finite set of groups, then the *graph product* of Γ and K, denoted Γ^K , is the quotient of the free product $*_{i \in [m]} \Gamma_i$ by all the relations $[\gamma_i, \gamma_j] = 1$, where $\gamma_i \in \Gamma_i, \gamma_j \in \Gamma_j$ and i and j are adjacent vertices of K. Note that Γ^K is the fundamental group of $X = (B\Gamma, *)^K$, where $B\Gamma = \{B\Gamma_1, ..., B\Gamma_m\}$ and * is a set of one-point subcomplexes. Note that by [34, Theorem 1.1], if K is a flag complex and each Γ_i is a discrete group, then $(B\Gamma, *)^K$ is a model for a $K(\Gamma^K, 1)$.

Theorem 3.15 Let \mathbb{F} and \mathcal{D} be skew fields and let K be a finite simplicial flag complex. Let $G = \mathbf{\Gamma}^K$ be a graph product of discrete groups such that there is an agrarian map $\mathbb{F}G \to \mathcal{D}$. If $H_{\bullet}(\Gamma; \mathcal{D}) = 0$ for each $\Gamma \in \mathbf{\Gamma}$, then

$$H_p^{\mathcal{D}}(\mathbf{\Gamma}) \cong \widetilde{H}_{p-1}(K; \mathcal{D}).$$

In particular, $b_p(G; \mathcal{D}) = b_{p-1}(K; \mathbb{F})$.

Proof Let $X = (B\Gamma, *)^K$ and let P be the poset whose elements are (possibly empty) simplices of K. Note that P defines a poset of spaces $\{X_{\sigma}\}_{\sigma \in P}$, where X_{\varnothing} is a single vertex. For each $\varnothing \neq \sigma \in P$, the group Γ^J is a direct product of groups with vanishing \mathcal{D} -homology, and therefore $H^n(\Gamma^J; \mathcal{D}) = H^n(X_J; \mathcal{D}) = 0$ by the Künneth formula. Take the spectral sequence of Lemma 2.3 with the coefficient system $\sigma \mapsto H_q(X_{\min(\sigma)}; \mathcal{D})$. All of these coefficient system cohomology groups vanish except when $\sigma = \emptyset$ and q = 0 and therefore

$$H^p(X; \mathcal{D}) = H^p(\operatorname{Flag}(\mathsf{P}), \operatorname{Flag}(\mathsf{P} > \emptyset); \mathcal{D}) \cong H^{p-1}(K; \mathcal{D})$$

by Lemma 2.3. The last isomorphism follows from the fact that P is isomorphic to the cone on the barycentric subdivision of K, where the cone point corresponds to the empty simplex \emptyset .

Remark 3.16 There are certain situations in which one can easily deduce the fact that a graph product has an agrarian embedding. For example, if each group $\Gamma \in \Gamma^{K}$ is ordered, then so is Γ^{K} by a result of Chiswell [10]. Thus, $\mathbb{F}\Gamma^{K}$ embeds into its Mal'cev–Neumann completion [28, 29].

Similarly, if each $\Gamma \in \Gamma^{K}$ is RFRS, then it is possible to show that Γ^{K} is also RFRS and therefore has a Hughes-free division ring $\mathcal{D}_{\mathbb{F}\Gamma^{K}}$. We provide a sketch of an argument here, which proceeds by induction on the number of vertices in *K*. If *K* has one vertex, then the claim is trivial. Suppose now that *K* has more than one vertex. If the 1-skeleton of *K* is a complete graph, then the claim is again trivial since Γ^{K} is a direct product of RFRS groups and therefore RFRS. If *K* is not a complete graph, then there is some vertex $v \in K$ such that $st(v) \neq K$ and we obtain a splitting $\Gamma^{K} \cong \Gamma^{st(v)} *_{\Gamma^{lk(v)}} \Gamma^{K \times v}$. By induction, both $\Gamma^{st(v)}$ and $\Gamma^{K \times v}$ are RFRS and by a result of Koberda and Suciu, the amalgam is also RFRS [22, Theorem 1.3]. Note that the Koberda–Suciu result is a combination theorem for a related class of groups called RFR *p* groups, but their proof is easily adapted to the RFRS case; we refer the interested reader to their paper for more details.

Thanks to the work of Okun and Schreve [30], we have the following corollary which holds, in particular, for RAAGs.

Corollary 3.17 Let \mathbb{F} and \mathcal{D} be skew fields and let K be a finite simplicial flag complex. Let $G = \Gamma^K$ be a graph product with an agrarian map $\mathbb{F}G \to \mathcal{D}$. If each $\Gamma \in \Gamma$ is residually finite, \mathcal{D} -acyclic, and \mathbb{F} - ℓ^2 -acyclic, then

$$b_n^{\mathcal{D}}(G) = \lim_{i \to \infty} \frac{b_n(G_i; \mathbb{F}_p)}{[G:G_i]}$$

for any residual chain $\Gamma = \Gamma_0 \bowtie G_1 \bowtie G_2 \bowtie \cdots$ of finite index normal subgroups.

Proof Okun and Schreve [30, Theorem 5.1] showed that the right-hand side of the above equation is independent of the choice of residual chain and equal to $\tilde{b}_{p-1}(K; \mathbb{F})$, which equals the left-hand side by Theorem 3.15.

Remark 3.18 Let K be a simplicial complex, $X = (B\Gamma, *)^K$, and let \widetilde{X} denote the universal cover. The above arguments apply equally well for computing $b_p^{\mathcal{D}}(\widetilde{X})$ and $b_p^{(2)}(X; \mathbb{F})$. In both cases they are equal to $\widetilde{b}_{p-1}(K; \mathbb{F})$ whenever they are defined. In the case when K is not a flag complex, X is not aspherical.

3.5 Artin groups

Let A be a residually finite Artin group and suppose the K(A, 1) conjecture holds for A. Then there is a contractible complex D_A —the *Deligne complex of A*—with stabilisers the maximal parabolic subgroups of A admitting a strict fundamental domain Q_A . In [30, Section 4 and Theorem 5.2] the authors compute $b_p^{(2)}(A; \mathbb{F}) = \tilde{b}_{p-1}(\partial Q_A; \mathbb{F})$ independently of a choice of residual chain. Here ∂Q is the subcomplex of Q with non-trivial stabilisers.

Theorem 3.19 Let \mathbb{F} be a field. Let A be a residually finite Artin group. Suppose the K(A, 1) conjecture holds for A. If $\mathbb{F}A \to \mathcal{D}$ is an agrarian map, then

$$b_p^{(2)}(A; \mathbb{F}) = b_p^{\mathcal{D}}(A; \mathbb{F}) = \tilde{b}_{p-1}(\partial Q_A; \mathbb{F}).$$

Proof In the action of A on D_A , the non-trivial stabilisers have a central \mathbb{Z} subgroup and so they are \mathcal{D} -acyclic and have vanishing \mathbb{F} -homology growth. We take a poset of spaces \mathcal{X} over the (barycentric subdivision of the) strict fundamental domain Q of A, where we assign a classifying space BA_{σ} to each $\sigma \in Q$. Now, we apply Remark 3.9 and Remark 3.5.

Remark 3.20 The above argument applies to residually finite groups G acting on a contractible complex with strict fundamental domain and \mathcal{D} -acyclic stabilisers—whenever G admits an agrarian embedding $\mathbb{F}G \to \mathcal{D}$.

3.6 Complements of hyperplane arrangements

Let \mathcal{A} be a collection of affine hyperplanes in \mathbb{C}^k and let $\Sigma(\mathcal{A})$ denote their union. Let $M(\mathcal{A}) := \mathbb{C}^k \setminus \Sigma(\mathcal{A})$. The *rank* of $M(\mathcal{A})$ is the maximum codimension *n* of any nonempty intersection of hyperplanes in \mathcal{A} . By [12, Proposition 2.1] the ordinary Betti numbers satisfy $\tilde{b}_p(M(\mathcal{A}); \mathbb{F}) = 0$ except possibly when p = n.

Theorem 3.21 Let \mathbb{F} be a skew field, \mathcal{A} be an affine hyperplane arrangement in \mathbb{C}^k of rank n, and let $\Gamma := \pi_1 M(A)$.

- (1) If Γ is residually finite, then $b_p^{(2)}(M(\mathcal{A}); \mathbb{F}) = \tilde{b}_p(M(\mathcal{A}); \mathbb{F})$ which equals zero except possibly when p = n.
- (2) If $\alpha : \mathbb{F}\Gamma \to \mathcal{D}$ is an agrarian map, then $b_p^{\mathcal{D}}(M(\mathcal{A})) = \widetilde{b}_p(M(\mathcal{A}); \mathbb{F})$ which equals zero except possibly when p = n.

We will need a lemma.

Lemma 3.22 Let \mathbb{F} be a skew field, \mathcal{A} be a non-empty central affine hyperplane arrangement in \mathbb{C}^k of rank n, and let $\Gamma := \pi_1 M(A)$.

- (1) If Γ is residually finite, then $b_p^{(2)}(M(\mathcal{A}); \mathbb{F}) = 0$ for all *n* independently of a choice of residual chain.
- (2) If $\alpha : \mathbb{F}\Gamma \to \mathcal{D}$ is an agrarian map, then $b_p^{\mathcal{D}}(M(\mathcal{A})) = 0$ for all n.

Proof In this case we have $M(A) = S^1 \times B$ where $B = M(A)/S^1$ by [12, Proof of Lemma 5.2]. Both results are easy applications of the Künneth formula and vanishing of the relevant invariant for $\mathbb{Z} = \pi_1 S^1$. We spell out the details in the first case to highlight the independence of the residual chain.

Observe that every finite cover M_i of $M(\mathcal{A})$ can be written as $B_i \times S^1$ such that S^1 has *m* one cells. We have the index of the cover $|M : M_i| = m|B : B_i|$. Now, we compute via the Künneth formula that

$$b_p^{(2)}(M, (M_i); \mathbb{F}) = \lim_{i \to \infty} \frac{b_p(M_i) + b_{p-1}(M_i)}{|M : M_i|}$$
$$= \lim_{i, m \to \infty} \frac{b_p(M_i) + b_{p-1}(M_i)}{m|B : B_i|}$$
$$= 0.$$

Proof of Theorem 3.21 By [12, Section 3] there is a cover \mathcal{U} of $M(\mathcal{A})$ by central subarrangements U_{σ} such that $\pi_1 U_{\sigma} \to \Gamma$ is injective and the nerve $N(\mathcal{U})$ is contractible. There is also a cover of a deleted neighbourhood of $\Sigma(A)$, denoted \mathcal{U}_{sing} , such that $H_p(N(\mathcal{U}), N(\mathcal{U}_{sing}))$ is concentrated in degree *n*. It follows from Lemma 3.22 that either the homology gradients or the \mathcal{D} -Betti numbers vanish. In particular, by Remarks 3.5 and 3.9 the cover \mathcal{U} is confident and the results follow.

4 A lower bound for homology gradients

Let *R* be a ring, let \mathcal{D} be a skew-field, and let $\varphi \colon R \to \mathcal{D}$ be a ring homomorphism. There is a rank function $\operatorname{rk}_{\mathcal{D},\varphi} \colon \operatorname{Mat}(R) \to \mathbb{R}_{\geq 0}$ defined by $\operatorname{rk}_{\mathcal{D},\varphi} A = \operatorname{rk} \varphi_* A$, where $\varphi_* A$ is the matrix obtained by applying the homomorphism φ to every entry of *A* and rk $\varphi_* A$ is the rank of $\varphi_* A$ as a matrix over \mathcal{D} . If $\varphi \colon R \hookrightarrow \mathcal{D}$ is an epic embedding (i.e. $\varphi(R)$ generates \mathcal{D} as a skew-field) and $\operatorname{rk}_{\mathcal{D},\varphi} \ge \operatorname{rk}_{\mathcal{E},\psi}$ for every skew-field \mathcal{E} and every ring homomorphism $\psi \colon R \to \mathcal{E}$, then \mathcal{D} is said to be a *universal division ring of fractions* for *R*. If \mathcal{D} is a universal division ring of fractions for *R*, it is then unique up to *R*-isomorphism [11, Theorem 4.4.1].

Theorem 4.1 (Jaikin-Zapirain [19, Corollary 1.3]) Let G be a residually (locally indicable and amenable) group and let \mathbb{F} be a skew-field. Then the Hughes-free division ring $\mathcal{D}_{\mathbb{F}G}$ exists and is the universal division ring of fractions of $\mathbb{F}G$.

We note that Theorem 4.1 holds for RFRS groups, since they are residually poly- \mathbb{Z} (see, e.g., [19, Proposition 4.4]). The main theorem of this section will follow quickly from the observation that $\mathcal{D}_{\mathbb{F}G}$ -Betti numbers scale under taking finite index subgroups and another result of Jaikin-Zapirain ([19, Corollary 1.6]), which he states for ℓ^2 -Betti numbers of CW complexes but also holds for agrarian homology. For the convenience of the reader, we reproduce a proof in the agrarian setting here.

Theorem 4.2 (Jaikin-Zapirain) Let \mathbb{F} be a skew-field and suppose that G is a group of type $\mathsf{FP}_{n+1}(\mathbb{F})$ for some $n \in \mathbb{N}$ such that $\mathcal{D}_{\mathbb{F}G}$ exists and is the universal division ring of fractions of $\mathbb{F}G$. Then $b_m^{\mathcal{D}_{\mathbb{F}G}}(G) \leq b_m(G; \mathbb{F})$ for all $m \leq n$.

Proof The embedding $\iota \colon \mathbb{F}G \hookrightarrow \mathcal{D}_{\mathbb{F}G}$ and the augmentation map $\alpha \colon \mathbb{F}G \to \mathbb{F}$ induce rank functions on Mat($\mathbb{F}G$) which we denote by rk_G and $\mathrm{rk}_{\mathbb{F}}$, respectively. By universality, $\mathrm{rk}_G \geq \mathrm{rk}_{\mathbb{F}}$.

Let $C_{\bullet} \to \mathbb{F}$ be a free-resolution of the trivial $\mathbb{F}G$ -module \mathbb{F} such that C_m is finitely generated for all $m \leq n + 1$. For all $m \leq n + 1$, let d_m be an integer such that $C_m \cong \mathbb{F}G^{d_m}$ and view the boundary maps $\partial_m \colon C_m \to C_{m-1}$ as matrices over $\mathbb{F}G$. The homologies we are interested in are $H_m(\mathcal{D}_{\mathbb{F}G} \otimes_{\mathbb{F}G} C_{\bullet})$ and $H_m(k \otimes_{\mathbb{F}G} C_{\bullet})$ and note that the differentials $\mathcal{D}_{\mathbb{F}G} \otimes \partial_m$ and $k \otimes \partial_m$ correspond to the matrices $\iota_* \partial_m$ and $\alpha_* \partial_m$ under the identifications $\mathcal{D}_{\mathbb{F}G} \otimes_{\mathbb{F}G} C_m \cong \mathcal{D}^{d_m}_{\mathbb{F}G}$ and $\mathbb{F} \otimes_{\mathbb{F}G} C_m \cong \mathbb{F}^{d_m}$. Therefore

$$b_m^{\mathcal{D}_{\mathbb{F}^G}}(G) = d_m - \mathrm{rk}_G \,\partial_m - \mathrm{rk}_G \,\partial_{m+1}$$
$$\leq d_m - \mathrm{rk}_{\mathbb{F}} \,\partial_m - \mathrm{rk}_{\mathbb{F}} \,\partial_{m+1}$$
$$= b_m(G; \,\mathbb{F})$$

As a consequence, we obtain that agrarian Betti numbers bound homology gradients from below.

Theorem 4.3 Let \mathbb{F} be a skew-field and let G be a group of type $\mathsf{FP}_{n+1}(\mathbb{F})$ such that $\mathcal{D}_{\mathbb{F}G}$ exists and is the universal division ring of fractions of $\mathbb{F}G$. If $H \leq G$ is any subgroup of finite index, then

$$b_m^{\mathcal{D}_{\mathbb{F}G}}(G) \le \frac{b_m(H;\mathbb{F})}{[G:H]}$$

for all $m \leq n$. In particular, if G is residually finite and $(G_i)_{i \in \mathbb{N}}$ is a residual chain of finite-index subgroups, then $b_m^{\mathcal{D}_{\mathbb{F}^G}}(G) \leq b_m^{(2)}(G, (G_i); \mathbb{F})$ for all $m \leq n$.

Proof By [15, Lemma 6.3], $[G : H] \cdot b_m^{\mathcal{D}_{\mathbb{F}G}}(G) = b_m^{\mathcal{D}_{\mathbb{F}H}}(H)$, and by Theorem 4.2. The second claim is an immediate consequence of the first.

5 Applications to fibring

A group *G* is *algebraically fibred* (or, simply, *fibred*) if it admits a nontrivial homomorphism $G \to \mathbb{Z}$ with finitely generated kernel. More generally, if \mathcal{P} is a finiteness property (e.g. type F_n or $FP_n(R)$ for some ring *R*, see Sect. 2.1 for definitions), we say that *G* is \mathcal{P} -*fibred* if there is a nontrivial homomorphism $G \to \mathbb{Z}$ with kernel of type \mathcal{P} .

Theorem 5.1 Let \mathbb{F} be a skew-field and let G be a RFRS group of type $\mathsf{FP}_{n+1}(\mathbb{F})$. If G is not virtually $\mathsf{FP}_n(\mathbb{F})$ -fibred, then for every residual chain of finite-index subgroups $(G_i)_{i \in \mathbb{N}}$, we have $b_m^{(2)}(G, (G_i); \mathbb{F}) > 0$ for some $m \leq n$.

Proof By [15] a RFRS group G is virtually $\mathsf{FP}_n(\mathbb{F})$ -fibred if and only if $b_i^{\mathcal{D}_{\mathbb{F}G}}(G) = 0$ for every $i \leq n$. Since G is not virtually $\mathsf{FP}_n(\mathbb{F})$ -fibred, we have $b_m^{\mathcal{D}_{\mathbb{F}G}}(G) > 0$ for some $m \leq n$. The result now follows from Theorem 4.3.

The authors thank Robert Kropholler for communicating to us the following question due to Matthew Zaremsky: If a RAAG A_L is virtually F_n -fibred, is it F_n -fibred? We are able to answer the analogous homological question over skew fields, \mathbb{Z} , and \mathbb{Z}/m for $m \in \mathbb{N}_{>1}$.

Theorem 5.2 Let *L* be a finite flag complex and *R* be either a skew field \mathbb{F} , \mathbb{Z} , or \mathbb{Z}/m for some $m \in \mathbb{N}_{>1}$. Then the RAAG A_L is virtually $\mathsf{FP}_n(R)$ -fibred if and only if it is $\mathsf{FP}_n(R)$ -fibred.

Proof We begin with the case $R = \mathbb{F}$. Let A_L be virtually $\mathsf{FP}_n(\mathbb{F})$ -fibred. By [15], a RFRS group *G* is virtually $\mathsf{FP}_n(\mathbb{F})$ -fibred if and only if $b_i^{\mathcal{D}_{\mathbb{F}G}}(G) = 0$ for every $i \leq n$. In particular this applies in the case $G = A_L$. By Theorem 3.15, $b_i^{\mathcal{D}_{\mathbb{F}A_L}}(A_L) = 0$ for every $i \leq n$ implies that $\tilde{b}_i(L; \mathbb{F}) = 0$ for every $i \leq n - 1$. By [5, Main Theorem], the Bestvina–Brady group is of type $\mathsf{FP}_n(\mathbb{F})$ and therefore A_L is $\mathsf{FP}_n(\mathbb{F})$ -fibred.

Now suppose $R = \mathbb{Z}/m$ for some $m \in \mathbb{N}_{>1}$ and suppose that A_L is virtually $\operatorname{FP}_n(\mathbb{Z}/m)$ -fibred. If p is a prime factor of m, then there is a ring homomorphism $\mathbb{Z}/m \to \mathbb{Z}/p = \mathbb{F}_p$ and therefore \mathbb{F}_p is a \mathbb{Z}/m -algebra. Thus, A_L is virtually $\operatorname{FP}_n(\mathbb{F}_p)$ -fibred. Therefore $b_i^{\mathcal{D}_{\mathbb{F}_pA_L}}(A_L) = 0$ for all $i \leq n$ and thus $\tilde{b}_i(L; \mathbb{F}_p) = 0$ for all $i \leq n-1$ for all $i \leq n-1$ by Theorem 3.15. Then $\tilde{b}_i(L; \mathbb{Z}/m) = 0$ for all $i \leq n-1$, so BB_L is of type $\operatorname{FP}_n(\mathbb{Z}/m)$.

Finally, suppose A_L is virtually FP_n -fibred. In particular, A_L is virtually $\mathsf{FP}_n(\mathbb{F})$ -fibred for every skew field \mathbb{F} , which implies that L is (n-1)-acyclic over every field by Theorem 3.15. Therefore L is (n-1)-acyclic over \mathbb{Z} , which implies that BB_L is of type FP_n .

Remark 5.3 In the case $\mathbb{F} = \mathbb{Q}$, Theorem 5.2 could have been deduced from previous work since the ℓ^2 -Betti numbers of RAAGs were computed by Davis and Leary in [13] and it is well known that a virtual FP_n(\mathbb{Q})-fibring implies the vanishing of ℓ^2 -Betti numbers in dimensions $\leq n$.

Since a finitely presented group of type FP_n is of type F_n we can reduce Zaremsky's question to one remaining case.

Question 5.4 (Zaremsky) Let L be a finite flag complex. If A_L is virtually F_2 -fibred, then is it F_2 -fibred?

We can also give examples of RAAGs that show that [15, Theorem A] does not hold when \mathbb{Q} is replaced by \mathbb{Z} . In other words, the vanishing of ℓ^2 -Betti numbers of RFRS groups does not detect virtual FP_n-fibrations.

Proposition 5.5 Let *p* be a prime. There are RAAGs that are ℓ^2 -acyclic but that do not virtually $FP_2(\mathbb{F}_p)$ -fibre. In particular, these RAAGs do not virtually FP_2 -fibre.

Proof Let *L* be a Q-acyclic flag complex that has non-trivial \mathbb{F}_p -homology in dimension 1, e.g. we may take *L* to be a flag triangulation of M(1, p) the Moore space with homology $\widetilde{H}_n(M(1, p); \mathbb{Z}) = 0$ unless n = 1, in which case it is isomorphic to \mathbb{Z}/p . Then A_L is ℓ^2 -acyclic by [13] (or Theorem 3.15) but it is not $\mathcal{D}_{\mathbb{F}_pA_L}$ -acyclic by Theorem 3.15. By [15, Theorem 6.6], A_L does not virtually $\mathsf{FP}_2(\mathbb{F}_p)$ -fibre and in particular does not virtually FP_n -fibre.

In contrast to this result, Kielak showed that if G is RFRS, of cohomological dimension at most 2, and ℓ^2 -acyclic, then G is virtually FP₂-fibred [21, Theorem 5.4].

The next application has to do with the following general question: If *G* fibres in two different ways, so that $G \cong K_1 \rtimes \mathbb{Z} \cong K_2 \rtimes \mathbb{Z}$ with K_1 and K_2 finitely generated, then what properties do K_1 and K_2 share? For example if *G* is a free-bycyclic (resp. surface-by-cyclic) group and $G \cong K \rtimes \mathbb{Z}$, then *K* is necessarily a free (resp. surface) group. We thank Ismael Morales for bringing the following question to our attention: if $G \cong K_1 \rtimes \mathbb{Z} \cong K_2 \rtimes \mathbb{Z}$ with K_1 and K_2 finitely generated, then is $b_1^{(2)}(K_1) = 0$ if and only if $b_1^{(2)}(K_2) = 0$? We prove this is the case for RAAGs, and obtain a similar result for higher ℓ^2 -Betti numbers and other agrarian invariants.

Theorem 5.6 Let $\varphi_0, \varphi_1 \colon A_L \to \mathbb{Z}$ be epimorphisms such that $BB_L^{\varphi_0}$ and $BB_L^{\varphi_1}$ are of type $\mathsf{FP}_n(\mathbb{F})$. If $\mathbb{F}BB_L^{\varphi_i} \hookrightarrow \mathcal{D}_i$ is an agrarian embedding for i = 0, 1, then $BB_L^{\varphi_0}$ is \mathcal{D}_0 -acyclic in dimensions $\leq n$ if and only if $BB_L^{\varphi_1}$ is \mathcal{D}_1 -acyclic in dimensions $\leq n$.

Before proving Theorem 5.6, we need a technical lemma. First we fix some notation. If $\sigma_1 = [e_1, \ldots, e_m]$ and $\sigma_2 = [f_1, \ldots, f_m]$ are ordered simplices in a simplicial complex *L* such that $\sigma_1 \cup \sigma_2$ is a simplex (equivalently, if $\sigma_1 \in \text{lk}(\sigma_2)$), then $\sigma_1 \cup \sigma_2$ always denotes the *ordered* simplex $[e_1, \ldots, e_m, f_1, \ldots, f_n]$. Moreover, if $\tau = \alpha_1 \sigma_1 + \cdots + \alpha_n \sigma_n$ is a formal linear combination of simplices σ_i (with coefficients α_i in some fixed skew field) such that $\sigma \cup \sigma_i \in L$ for every *i*, then $\sigma \cup \tau$ denotes the formal linear combination

$$\alpha_1 \sigma \cup \sigma_1 + \cdots + \alpha_n \sigma \cup \sigma_n.$$

If $\varphi: A_L \to \mathbb{Z}$ is a homomorphism, recall that L^a is the subcomplex of L spanned by the vertices $v \in L$ such that $\varphi(v) \neq 0$. We will write $lk_L(\sigma)$ (resp. $lk_{L^a}(\sigma)$) for the link of a simplex σ in L (resp. L^a).

Lemma 5.7 Let BB_L^{φ} be of type $FP_n(\mathbb{F})$ and let v be a dead vertex of L. Then every simplicial (n - 1)-cycle of $lk_L(v)$ is homologous to a cycle in $lk_{L^a}(v)$.

Proof Let $\sigma = \alpha_1 \sigma_1 + \cdots + \alpha_k \sigma_k$ be a simplicial (n-1)-cycle in $\mathbb{l}_L(v)$, where each σ_i is an ordered (n-1)-simplex of $\mathbb{l}_L(v)$ and $\alpha_i \in \mathbb{F}$ for each *i*. By induction on $m \ge 0$, we will show that the simplices σ_i can be replaced with (n-1)-simplices having at least *m* living vertices such that the resulting chain is a cycle homologous to σ . The lemma follows from the m = n case.

For the base case, suppose that σ_i is a simplex with no living vertices. Then $\{v\} \cup \sigma_i$ is a dead *n*-simplex and therefore $lk_{L^a}(\{v\} \cup \sigma_i)$ is (-1)-connected (see Remark 3.13), i.e. it is nonempty. Thus, there is a living vertex *u* such that $\{u\} \cup \sigma_i \subseteq lk_L(v)$. Since

$$\partial(\{u\}\cup\sigma_i)=\sigma_i-\{u\}\cup\partial\sigma_i,$$

where $\{u\} \cup \partial \sigma_i$ is a linear combination of (n - 1)-simplices with one living vertex, we can replace σ_i with $\{u\} \cup \partial \sigma_i$ in σ . Hence, we assume that the linear combination $\alpha_1 \sigma_1 + \cdots + \alpha_k \sigma_k$ only involves simplices with at least one living vertex.

Assume that $\alpha_1 \sigma_1 + \cdots + \alpha_k \sigma_k$ only involves simplices with at least $m \ge 1$ living vertices for some m < n and let σ_i be a simplex with exactly *m* living vertices.

Let $\lambda \subseteq \sigma_i$ be the dead (n - m - 1)-face of σ_i and let $\sigma_i = \sigma_{i_1}, \ldots, \sigma_{i_l}$ be the simplices among $\{\sigma_1, \ldots, \sigma_k\}$ containing λ as a face. For each $j \in \{1, \ldots, l\}$, write $\sigma_{i_j} = \varepsilon_j \lambda \cup \tau_j$, where τ_j is a living (m - 1)-simplex of $lk_L(v)$ and $\varepsilon_j \in \{\pm 1\}$. Then

$$\begin{split} \partial \left(\sum_{j=1}^{l} \alpha_{i_j} \sigma_{i_j} \right) &= \sum_{j=1}^{l} \alpha_{i_j} \varepsilon_j (\partial \lambda \cup \tau_j + (-1)^{n-m} \lambda \cup \partial \tau_j) \\ &= \left(\sum_{j=1}^{l} \alpha_{i_j} \varepsilon_j \partial \lambda \cup \tau_j \right) + (-1)^{n-m} \lambda \cup \partial \left(\sum_{j=1}^{l} \alpha_{i_j} \varepsilon_j \tau_j \right) \\ &= 0, \end{split}$$

since $\partial \sigma = 0$ and the simplices σ_{i_j} are the only simplices among $\{\sigma_1, \ldots, \sigma_k\}$ containing λ as a face. Thus,

$$\lambda \cup \partial \left(\sum_{j=1}^{l} \alpha_{i_j} \varepsilon_j \tau_j \right) = 0,$$

whence we conclude that $\sum_{j=1}^{l} \alpha_{i_j} \varepsilon_j \tau_j$ is an (m-1)-cycle in $lk_{L^a}(\{v\} \cup \lambda)$. But $\{v\} \cup \lambda$ is a dead (n-m)-simplex and therefore $lk_{L^a}(\{v\} \cup \lambda)$ is (m-1)-connected. Hence, $\sum_{j=1}^{l} \alpha_{i_j} \varepsilon_j \tau_j = \partial \psi$ for some living *m*-chain ψ in $lk_{L^a}(\{v\} \cup \lambda)$. Then

$$\partial(\lambda \cup \psi) = \partial\lambda \cup \psi + (-1)^{n-m}\lambda \cup \left(\sum_{j=1}^{l} \alpha_{i_j}\varepsilon_j\tau_j\right)$$
$$= \partial\lambda \cup \psi + (-1)^{n-m}\sum_{j=1}^{l} \alpha_{i_j}\sigma_{i_j}.$$

The chain $\partial \lambda \cup \psi$ is a linear combination of simplices with m + 1 living vertices. We can therefore replace $\sum_{j=1}^{l} \alpha_{ij} \sigma_{ij}$ with $\pm \partial \lambda \cup \psi$ and assume that σ is a linear combination of simplices each with at least m + 1 living vertices.

Proof of Theorem 5.6 Suppose that $b_p^{\mathcal{D}_0}(BB_L^{\varphi_0}) > 0$ for some $p \leq n$. By Theorem 3.12, there is a vertex v of L such that $\varphi_0(v) \neq 0$ and

$$\widetilde{b}_{p-1}(\operatorname{lk}(v); \mathbb{F}) > 0.$$

Hence, there is a simplicial (p-1)-cycle σ in lk(v) that is not a boundary. If $\varphi_1(v) = 0$, then, by Lemma 5.7, σ is homologous to a cycle in lk_{L^a}(v) where L^a denotes the living link with respect to φ_1 . Thus $\widetilde{H}_{p-1}(\text{lk}_{L^a}(v); \mathbb{F}) \neq 0$. But lk_{L^a}(v) is (n-1)-connected over \mathbb{F} , so we must have $\varphi_1(v) \neq 0$, and therefore $b_p^{\mathcal{D}_1}(BB_L^{\varphi_1}) > 0$ by Theorem 3.12.

We highlight the following immediate corollary.

Corollary 5.8 *Either all the* $\operatorname{FP}_n(\mathbb{F})$ *-fibres of* A_L *are virtually* $\operatorname{FP}_n(\mathbb{F})$ *-fibred or none of them are. In particular, either all of* A_L *'s fibres virtually fibre or none of them do.*

Proof This follows from Theorem 5.6 and the fact that being $\mathcal{D}_{\mathbb{F}BB_L^{\varphi}}$ -acyclic in dimensions $\leq n$ is equivalent to virtually fibring with kernel of type $\mathsf{FP}_n(\mathbb{F})$ [15, Theorem 6.6].

6 Amenable category and minimal volume entropy

In this section we will relate $\mathcal{D}_{\mathbb{F}G}$ -Betti numbers with amenable category and minimal volume entropy.

Let *X* be a path-connected space with fundamental group *G*. A (not necessarily path-connected) subset *U* of *X* is an *amenable subspace* if $\pi_1(U, x) \rightarrow \pi(X, x)$ has amenable image for all $x \in U$. The *amenable category*, denoted $\operatorname{cat}_{AMN}X$, is the minimal $n \in \mathbb{N}$ for which there exists an open cover of *X* by n+1 amenable subspaces. If no such cover exists we set $\operatorname{cat}_{AMN}X = \infty$. The definition of amenable category has been extracted from [9] and [23]. Note that we normalise the invariant as in the second paper. Also note that often in the literature the multiplicity of the cover is considered instead, however, the two definitions turn out to be equivalent for CW complexes [9, Remark 3.13]

Proposition 6.1 Let \mathbb{F} be a skew field. Let G be residually finite of type F , and suppose $\mathcal{D}_{\mathbb{F}G}$ exists. If $\mathsf{cat}_{\mathcal{AMN}G} = k$. Then, $b_p^{\mathcal{D}_{\mathbb{F}G}}(G) = b_p^{(2)}(G, (G_n); \mathbb{F}) = 0$ for $p \ge k-1$ and every residual chain (G_n) .

Proof Let X be a finite model for a K(G, 1). As explained in [17, Theorem 3.2] we may adapt the proof of [31, Theorem 1.6] to apply to k-dimensional aspherical simplicial complexes. In particular, for a residual chain $(G_n)_{n \in \mathbb{N}}$ we obtain a sequence of covers $X_n \to X$, such that the number of p-cells in X_n grows sublinearly in $[G : G_n]$. Since

$$b_p^{\mathcal{D}_{\mathbb{F}^G}}(X_n) = [G:G_n] \cdot b_p^{\mathcal{D}_{\mathbb{F}^G}}(G) \le |I_n(X_k)|$$

where $I_p(X_k)$ is the set of *p*-cells of X_n . But now, as *k* tends to infinity, the left hand side of the equation grows linearly, and the right hand side of the equation grows sublinearly. This is only possible if $b_n^{\mathcal{D}_{\mathbb{F}G}}(G) = 0$. The statement concerning $b_p^{(2)}(G, (G_n); \mathbb{F})$ is analogous.

Let X be a finite CW complex with a piecewise Riemannian metric g. Fix a basepoint x_0 in the universal cover \widetilde{X} and let \widetilde{g} be the pull-back metric. The *volume entropy* of (X, g) is

$$\operatorname{ent}(X,g) := \lim_{t \to \infty} \frac{1}{t} \operatorname{Vol}(B_{x_0}(t),\widetilde{g}).$$

The *minimal volume entropy* of *X* is

$$\omega(X) := \inf_{g} \operatorname{ent}(X, g) \operatorname{Vol}(X, g)^{1/\dim X}$$

where g varies over all piecewise Riemannian metrics. The invariant was originally defined for Riemannian manifolds in [16].

Suppose G is a group admitting a finite K(G, 1). The *minimal volume entropy* of G is

$$\omega(G) := \inf(\omega(X))$$

where *X* ranges over all finite models of a K(G, 1) such that dim X = gd(G).

There are few calculations of minimal volume entropy of groups which are not fundamental groups of aspherical manifolds in literature. To date there is the work of Babenko–Sabourau [2] on which computations for free-by-cyclic groups [6] and RAAGs [17, 23] have been completed.

We say G has uniformly uniform exponential growth if each subgroup either has uniform exponential growth bounded below by some constant $\omega_0 > 1$ or is virtually abelian. Note that this property is sometimes called *uniform uniform exponential* growth or locally uniform exponential growth.

Corollary 6.2 Let \mathbb{F} be a skew field. Let G be a residually finite group of type F , and suppose $\mathcal{D}_{\mathbb{F}G}$ exists. If G has uniformly uniform exponential growth and is not $\mathcal{D}_{\mathbb{F}G}$ -acyclic, then $\omega(G) > 0$.

Proof This follows from [17, Paragraph after Theorem 3.3] swapping out their use of [17, Theorem 3.3] for our Proposition 6.1. □

Corollary 6.3 Let \mathbb{F} be a skew field and let $\varphi \colon A_L \to \mathbb{Z}$ be an epimorphism. Suppose BB_L^{φ} is of type F. If $\bigoplus_{v \in L^2} \widetilde{H}_{p-1}(\operatorname{lk}(v); \mathbb{Z}) \neq 0$, then $\omega(BBL_L^{\varphi}) > 0$.

Proof This follows from Corollary 6.2 and the fact that a right-angled Artin group has strongly uniform exponential growth by [3].

We conjecture that the converse of the last corollary holds.

Conjecture 6.4 Let \mathbb{F} be a skew field and let $\varphi \colon A_L \to \mathbb{Z}$ be an epimorphism. Suppose BB_L^{φ} is of type F. If $\bigoplus_{v \in L^a} \widetilde{H}_{p-1}(\operatorname{lk}(v); \mathbb{Z}) = 0$, then $\omega(BBL_L^{\varphi}) = 0$.

Acknowledgements Sam P. Fisher is supported by the National Science and Engineering Research Council (NSERC) [ref. no. 567804-2022]. Sam Hughes received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 850930). The authors would like to thank Ismael Morales for asking a question that led to Theorem 5.6, as well as Robert Kropholler and Matt Zaremsky for communicating a question which inspired Theorem 5.2. The authors would like to thank Dawid Kielak for helpful conversations.

Data availibility Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Avramidi, G., Okun, B., Schreve, K.: Mod p and torsion homology growth in nonpositive curvature. Invent. Math. 226(3), 711–723 (2021)
- 2. Babenko, I., Sabourau, S.: Minimal volume entropy and fiber growth (2021). arXiv:2102.04551 [math.GT]
- 3. Baudisch, A.: Subgroups of semifree groups. Acta Math. Acad. Sci. Hungar. 38(1-4), 19-28 (1981)
- Bergeron, N., Linnell, P., Lück, W., Sauer, R.: On the growth of Betti numbers in *p*-adic analytic towers. Groups Geom. Dyn. 8(2), 311–329 (2014)
- Bestvina, M., Brady, N.: Morse theory and finiteness properties of groups. Invent. Math. 129(3), 445– 470 (1997)
- Bregman, C., Clay, M.: Minimal volume entropy of free-by-cyclic groups and 2-dimensional rightangled Artin groups. Math. Ann. 381(3–4), 1253–1281 (2021)
- Brown, K.S.: Cohomology of groups, Graduate Texts in Mathematics, vol. 87. Springer, New York (1994). Corrected reprint of the 1982 original
- Bux, K.-U., Gonzalez, C.: The Bestvina–Brady construction revisited: geometric computation of Σinvariants for right-angled Artin groups. J. Lond. Math. Soc. (2) 60(3), 793–801 (1999)
- Capovilla, P., Löh, C., Moraschini, M.: Amenable category and complexity. Algebr. Geom. Topol. 22(3), 1417–1459 (2022)
- 10. Chiswell, I.M.: Ordering graph products of groups. Int. J. Algebra Comput. 22(4), 1250037 (2012)
- Cohn, P.M.: Skew Fields: Theory of General Division Rings. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge (1995)
- Davis, M.W., Januszkiewicz, T., Leary, I.J.: The l²-cohomology of hyperplane complements. Groups Geom. Dyn. 1(3), 301–309 (2007)
- Davis, M.W., Leary, I.J.: The l²-cohomology of Artin groups. J. Lond. Math. Soc. (2) 68(2), 493–510 (2003)
- Davis, M.W., Okun, B.: Cohomology computations for Artin groups, Bestvina–Brady groups, and graph products. Groups Geom. Dyn. 6(3), 485–531 (2012)
- 15. Fisher, S.P.: Improved algebraic fibrings (2021). arXiv:2112.00397
- 16. Gromov, M.: Volume and bounded cohomology. Inst. Hautes Études Sci. Publ. Math. 6, 5-99 (1983)
- 17. Haulmark, M., Schreve, K.: Minimal volume entropy of RAAG's (2022). arXiv:2202.10405 [math.GR]
- 18. Henneke, F., Kielak, D.: Agrarian and L^2 -invariants. Fundam. Math. **255**(3), 255–287 (2021)
- Jaikin-Zapirain, A.: The universality of Hughes-free division rings. Selecta Math. (N.S.) 27(4), Paper No. 74, 33 (2021)
- Jaikin-Zapirain, A., López-Álvarez, D.: The strong Atiyah and Lück approximation conjectures for one-relator groups. Math. Ann. 376(3–4), 1741–1793 (2020)
- Kielak, D.: Residually finite rationally solvable groups and virtual fibring. J. Am. Math. Soc. 33(2), 451–486 (2020)
- 22. Koberda, T., Suciu, A.I.: Residually finite rationally p groups. Commun. Contemp. Math. **22**(3), 1950016 (2020)
- 23. Li, K.: Amenable covers of right-angled Artin groups (2022). arXiv:2204.01162 [math.GR]

- 24. Linnell, P.A.: Division rings and group von Neumann algebras. Forum Math. 5(6), 561–576 (1993)
- 25. Linnell, P., Lück, W., Sauer, R.: The limit of \mathbb{F}_p -Betti numbers of a tower of finite covers with amenable fundamental groups. Proc. Am. Math. Soc. **139**(2), 421–434 (2011)
- 26. Lück, W.: Approximating L^2 -invariants by their finite-dimensional analogues. Geom. Funct. Anal. **4**(4), 455–481 (1994)
- Lück, W.: Approximating L²-invariants by their classical counterparts. EMS Surv. Math. Sci. 3(2), 269–344 (2016)
- Mal'cev, A.I.: On the embedding of group algebras in division algebras. Doklady Akad. Nauk SSSR (N.S.) 60, 1499–1501 (1948)
- 29. Neumann, B.H.: On ordered division rings. Trans. Am. Math. Soc. 66, 202-252 (1949)
- 30. Okun, B., Schreve, K.: Torsion invariants of complexes of groups (2021). arXiv:2108.08892 [math.GR]
- Sauer, R.: Volume and homology growth of aspherical manifolds. Geom. Topol. 20(2), 1035–1059 (2016)
- 32. Schick, T.: Erratum: "Integrality of L²-Betti numbers". Math. Ann. **322**(2), 421–422 (2002)
- Schreve, K.: The strong Atiyah conjecture for virtually cocompact special groups. Math. Ann. 359(3– 4), 629–636 (2014)
- Stafa, M.: On the fundamental group of certain polyhedral products. J. Pure Appl. Algebra 219(6), 2279–2299 (2015)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.