

QUASIHOMOMORPHISMS TO REAL ALGEBRAIC GROUPS

SAMI DOUBA, FRANCESCO FOURNIER-FACIO, SAM HUGHES, AND SIMON MACHADO

ABSTRACT. A quasihomomorphism is a map that satisfies the homomorphism relation up to bounded error. Fujiwara and Kapovich proved a rigidity result for quasihomomorphisms taking values in discrete groups, showing that all quasihomomorphisms can be built from homomorphisms and sections of bounded central extensions. We study quasihomomorphisms with values in real linear algebraic groups, and prove an analogous rigidity theorem.

1. INTRODUCTION

Let G be a locally compact group. We say that a subset $D \subset G$ is *bounded* if D is contained in a compact subset of G . Let Γ be a group. A map $f: \Gamma \rightarrow G$ is a *quasihomomorphism* if there exists a bounded subset $D_f \subset G$ such that

$$f(xy) \in f(x)f(y)D_f \text{ for all } x, y \in \Gamma.$$

Two quasihomomorphisms $f, g: \Gamma \rightarrow G$ are *close* if there exists a bounded subset $D_{f,g} \subset G$ such that $f(x) \in g(x)D_{f,g}$ for all $x \in \Gamma$. Note that the notion of quasihomomorphism depends on the topology of G , but not on the topology of Γ , so we always view Γ as a discrete group.

Our main result is a rigidity theorem for quasihomomorphisms with values in a real linear¹ algebraic group, in the spirit of [FK16]. We define a *bounded modification* of a quasihomomorphism $f: \Gamma \rightarrow G$ to be a quasihomomorphism² $f': \Gamma' \rightarrow G'$, where $\Gamma' < \Gamma$ is a finite-index subgroup, $G' < G$ is a closed subgroup, and f' is close to $f|_{\Gamma'}$. If G has the structure of a real algebraic group, then we require moreover that G' be an algebraic subgroup of G . A subset X of a group G is *normal* if $g^{-1}Xg = X$ for all $g \in G$.

Theorem A (Main theorem, succinct form). *Let $f: \Gamma \rightarrow G$ be a quasihomomorphism, where G is a real algebraic group. Then there exists a bounded modification $f': \Gamma' \rightarrow G'$ such that $D_{f'}$ is contained in a bounded normal subset of G' .*

There is a rich theory of quasihomomorphisms when the target is an abelian group, thanks to their many appearances throughout mathematics: stable commutator length [Cal09], bounded cohomology [Fri17], one-dimensional dynamics [Ghy87], geometric group theory [BF02, Man08], knot theory [Mal04] and symplectic geometry [PR14]. Rigidity results for quasihomomorphisms can be traced back to work of Burger–Monod [BM99, BM02] on bounded cohomology of higher-rank lattices. This motivated Ozawa [Oza11] to introduce property (TTT) as a tool to prove rigidity of quasihomomorphisms with amenable target. Dumas [Dum24] has since proved that lattices in higher-rank simple algebraic groups over local fields satisfy this property. Recently, the work of Hrushovski [Hru22] showed that

Date: 4th March, 2026.

Mathematical subject classification 2020. 20C99, 20F65, 20G20, 20J05, 22E15.

¹In the sequel, the word “linear” will be implicit.

²Outside the case of abelian target, a map that is close to a quasihomomorphism need not be a quasihomomorphism (Example 3.5).

quasihomomorphisms play a central role in model theory, especially in relation to approximate subgroups.

The result in this direction that is most relevant to the present work is due to Fujiwara–Kapovich [FK16], who studied the case in which the target is discrete (so that bounded sets are finite). They proved that if $f: \Gamma \rightarrow G$ is a quasihomomorphism with discrete target, then there exists a bounded modification $f': \Gamma' \rightarrow G'$ such that $D_{f'}$ is *central* in G' . Therefore, up to passing to a finite-index subgroup and changing by a bounded amount, one can construct f from homomorphisms and sections of bounded central extensions. Accordingly, they call such quasihomomorphisms *constructible*.

In the non-discrete setting, one cannot expect such a strong rigidity statement, even for $G = \mathrm{SO}(2) \times \mathbb{R}^2$: see Proposition 6.1. As remarked by Hrushovski [Hru22, Section 5.17], the right generalisation of “central” is “normal” once one abandons the discrete world.

Remark. There appears to be some confusion in the literature about constructibility of quasihomomorphisms: [BV24] claims the existence of non-constructible quasihomomorphisms in the case where Γ is the fundamental group of a closed hyperbolic manifold and G is a non-abelian unipotent real algebraic group. Below, we exhibit a counterexample to [BV24, Theorem 4.28]³; see Example 3.6 and Remark 3.7 for a discussion.

Quasihomomorphisms with normal defect play a special role in the literature; indeed, they are the ones that appear in relation to model theory [Hru22]. It turns out that such quasihomomorphisms are very restricted, in the general case of a locally compact target [Hru22, Proposition C.3]. In our case we can push this even further, giving the following stronger version of Theorem A. Following Hrushovski [Hru22], we say that a simply connected abelian algebraic normal subgroup $A \triangleleft G$ is *rigid* if the homomorphism $G \rightarrow \mathrm{Aut}(A)$ induced by conjugacy has compact image.⁴ In this case, $A \cong \mathbb{R}^n$ for some $n \in \mathbb{N}$, and there is a Euclidean structure on A such that the action of G factors through $\mathrm{O}(n)$.

Theorem B (Main theorem, stronger form). *Let $f: \Gamma \rightarrow G$ be a quasihomomorphism, where G is a real algebraic group. Then there exists a bounded modification $f': \Gamma' \rightarrow G'$ with the following property:*

There exists a compact (algebraic) normal subgroup $C \triangleleft G'$ and a rigid abelian algebraic normal subgroup $A \triangleleft G'/C$ such that $D_{f'}$ is contained in the preimage of A in G' .

It follows that the composition $\Gamma' \xrightarrow{f'} G' \rightarrow (G'/C)/A$ is a homomorphism. Therefore, up to passing to a finite-index subgroup and changing by a bounded amount, one can build f from homomorphisms, sections of bounded *abelian* extensions, and sections of compact extensions. Since bounded abelian extensions are governed by bounded cohomology, we obtain the following corollary.

Corollary C. *Let Γ be a group with the property that for every finite-dimensional orthogonal representation π of Γ with no non-zero invariant vectors, the second bounded cohomology $H_b^2(\Gamma; \pi)$ vanishes. Then for every quasihomomorphism $f: \Gamma \rightarrow G$, where G is a real algebraic group, there exists a bounded modification $f': \Gamma' \rightarrow G'$, and a compact (algebraic) normal subgroup $C \triangleleft G'$ such that the composition $\Gamma' \xrightarrow{f'} G' \rightarrow G'/C$ is a quasihomomorphism with central defect.*

³Theorem 4.24 in the arXiv version

⁴Hrushovski also allows for \mathbb{Z}^n factors, but these do not occur in the algebraic setting.

If moreover the second bounded cohomology with trivial coefficients $H_b^2(\Gamma; \mathbb{R})$ vanishes, then $f': \Gamma' \rightarrow G'$ and $C \triangleleft G'$ can be chosen such that $\Gamma' \xrightarrow{f'} G' \rightarrow G'/C$ is a homomorphism.

The first statement of Corollary C applies to lattices in higher-rank simple groups, while the second statement applies only in the non-Hermitian case [MS04]. Moreover, the second statement applies to amenable groups [Joh72], certain lamplighters [Mon22, Bog25], Thompson's group F and other groups of dynamical origin such as compactly supported transformation groups of Euclidean spaces [Mon22, CFFLM25] and certain natural colimit groups arising in the context of homological stability [CFFLM25]. In the terminology of [FK16], the last conclusion is that the quasihomomorphism f' is an *almost homomorphism*.

Under stronger assumptions on the target group, we obtain severe restrictions on quasihomomorphisms, as demonstrated by the following result, which serves as a starting point for the proof of the main theorem.

Theorem D. *Let \mathbb{K} be a local field, let $G = \mathbf{G}(\mathbb{K})$, where \mathbf{G} is a connected adjoint simple \mathbb{K} -group, and let $f: \Gamma \rightarrow G$ be a quasihomomorphism. If $f(\Gamma)$ is unbounded in G and generates a Zariski-dense subgroup of G , then f is a homomorphism.*

Theorem D holds also for non-Archimedean local fields, in particular it holds for p -adic groups. However, in the final step of our proof of the main theorem, we need the fact that a real algebraic group with a compact Zariski-dense subgroup is itself compact (see the footnote in Section 5.D), hence our restriction to the setting of real algebraic groups.

A consequence of the proof of the main theorem is that the difference between *normal* and *central* defect is only due to the presence of compact quotients, as in Proposition 6.1. Therefore, excluding compact quotients, we obtain the following direct analogue of [FK16].

Theorem E. *Let $f: \Gamma \rightarrow G$ be a quasihomomorphism, where G is a real algebraic group. Suppose that the Zariski-closure in G of the group generated by $f(\Gamma)$ has no non-trivial compact algebraic quotients. Then there exists a bounded modification $f': \Gamma \rightarrow G'$ such that $D_{f'}$ is central in G' .*

Note that here we did not need to pass to a finite-index subgroup of Γ . More generally, in the proof of the main theorem, we need only pass to a finite-index subgroup to ensure that the Zariski-closure of the group generated by the image of the quasihomomorphism is Zariski-connected (Lemma 5.1).

We believe that the analogue of Theorem A should hold in the more general setting of real Lie groups. Together with the Gleason–Yamabe Theorem [Tao14] (using Remark 2.7), this would then give a rigidity result for quasihomomorphisms with connected locally compact target. In the opposite direction, we do not know whether to expect a positive answer to the following:

Question F. *Is there a quasihomomorphism $f: \Gamma \rightarrow G$, where G is a totally disconnected, locally compact group, for which Theorem A does not hold? What if one assumes $\Gamma = \mathbb{Z}$?*

Outline. We start with generalities on quasihomomorphisms in Section 2. Then in Section 3 we prove Theorem D, which serves as a starting point for the proof of the main theorem. In Section 4, we prove some technical results on real algebraic groups, most notably Proposition 4.3. In Section 5 we prove the main theorem in all its various versions: Theorem A, Theorem B, Corollary C and Theorem E. Finally, in Section 6, we prove Proposition 6.1, which shows that in our most general setting one cannot hope for a bounded modification with central defect.

Acknowledgements. SD and SH are supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC-2047/1 - 390685813. FFF is supported by the Herchel Smith Postdoctoral Fellowship Fund. SH is supported by a Humboldt Research Fellowship at Universität Bonn. Various stages of this research were conducted at the Forschungsinstitut für Mathematik (ETH Zürich); at the Isaac Newton Institute for Mathematical Sciences during the programme Operators, Graphs, Groups (EPSRC grant EP/Z000580/1s); and at the Simons Laufer Mathematical Sciences Institute in Berkeley, California, during the Spring 2026 semester (National Science Foundation Grant No. DMS-2424139); the authors acknowledge these institutions' hospitality and support.

2. GENERALITIES ON QUASIHOMOMORPHISMS

Notation 2.1. We work with right actions. Accordingly, conjugacy is denoted by $g^h = h^{-1}gh$, and commutators by $[g, h] = g^{-1}h^{-1}gh = g^{-1}g^h$.

Let K be a bounded set. If $x \in yK$, then we write $x \approx_K y$. Below we will introduce shorthand notations to specify what the compact set is: see Lemma 2.6 and the proof of Claim 5.8.

When H is a subgroup of the algebraic group G , we denote by $\overline{H}^{\text{Zar}}$ the Zariski-closure of H in G .

Definition 2.2. Let $f: \Gamma \rightarrow G$ be a map to a locally compact group G . The *defect set* of f is

$$D_f := \{f(y)^{-1}f(x)^{-1}f(xy) \mid x, y \in \Gamma\}.$$

We say that f is a *quasihomomorphism* if D_f is bounded. The *defect group* of f is the topological closure of the group generated by D_f and is denoted by Δ_f .

Using the notation above, f is a quasihomomorphism if $f(xy) \approx_K f(x)f(y)$ for some bounded set K . Similarly, f and g are close if $f(x) \approx_K g(x)$.

Lemma 2.3. [FK16, Section 2] *Let $f: \Gamma \rightarrow G$ be a function. Then*

$$D_f^{f(x)} \subset D_f^2 D_f^{-1}$$

for all $x \in \Gamma$.

Lemma 2.4. *Let $x \in \langle D_f \rangle$. Then $x^{f(\Gamma)}$ is bounded.*

Proof. By Lemma 2.3, we have

$$x^{f(\Gamma)} \subset (D_f^{f(\Gamma)} \cup (D_f^{-1})^{f(\Gamma)})^k \subset (D_f^2 D_f^{-1} \cup D_f D_f^{-2})^k. \quad \square$$

Proposition 2.5. *Suppose that G is an algebraic group, and that $f: \Gamma \rightarrow G$ is a quasihomomorphism. Suppose that $f(\Gamma)$ generates a Zariski-dense subgroup of G . Then $\overline{\Delta_f}^{\text{Zar}} = \overline{\langle D_f \rangle}^{\text{Zar}}$ is a normal subgroup of G .*

Proof. Lemma 2.3 implies that every element of $f(\Gamma)$ normalises $\langle D_f \rangle$, hence also Δ_f . It follows that the group generated by $f(\Gamma)$ normalises Δ_f , and hence also $\overline{\Delta_f}^{\text{Zar}}$. This is an algebraic condition, therefore by Zariski-density G normalises $\overline{\Delta_f}^{\text{Zar}}$, too. \square

Lemma 2.6. *Let $y_1, \dots, y_m, z_1, \dots, z_m \in \Gamma$ and let $f: \Gamma \rightarrow G$ be a quasihomomorphism. Then*

$$f(y_1 z_1^{-1} \cdots y_m z_m^{-1}) \approx_{K_m} f(y_1) f(z_1)^{-1} \cdots f(y_m) f(z_m)^{-1},$$

where K_m is the set $(D_f \cup D_f^{-1})$ to a power depending only on m .

Proof. For $m = 1$, first note that

$$f(1) = f(1 \cdot 1) \in f(1)f(1)D_f \quad \Rightarrow \quad f(1) \in D_f^{-1}.$$

Next,

$$f(1) = f(xx^{-1}) \in f(x)f(x^{-1})D_f \quad \Rightarrow \quad f(x)^{-1} \in f(x^{-1})D_f^2.$$

It follows that

$$f(yz^{-1}) \in f(y)f(z^{-1})D_f \subset f(y)f(z)^{-1}D_f^{-2}D_f.$$

So $K_1 = (D_f \cup D_f^{-1})^3$ works.

Suppose that the statement is true up to m , and let K_m be the corresponding bounded set. Then

$$\begin{aligned} f(y_1 z_1^{-1} \cdots y_{m+1} z_{m+1}^{-1}) &\in f(y_1 z_1^{-1})f(y_2 z_2^{-1} \cdots y_{m+1} z_{m+1}^{-1})D_f \\ &\subset f(y_1)f(z_1)^{-1}K_1 f(y_2)f(z_2)^{-1} \cdots f(y_{m+1})f(z_{m+1})^{-1}K_m D_f. \end{aligned}$$

Using Lemma 2.3, we can conjugate K_1 by appropriate elements in the image of f while keeping the error set a power of $(D_f \cup D_f^{-1})$. \square

Remark 2.7. Let $\pi: G \rightarrow Q$ be quotient with compact kernel C . If $f: \Gamma \rightarrow G$ is a quasimorphism, then $D_{\pi f} = \pi(D_f)$, hence πf is also a quasihomomorphism. Conversely, if $g: \Gamma \rightarrow Q$ is a quasihomomorphism, then and $\hat{g}: \Gamma \rightarrow G$ is a set-theoretic lift, then $D_{\hat{g}} \subset \pi^{-1}D_g$, hence \hat{g} is also a quasihomomorphism.

It is also easy to see that the conclusion of our main theorem (in its various forms: Theorem A, Theorem B and Theorem E) is preserved in passing from $f: \Gamma \rightarrow G$ to $\pi f: \Gamma \rightarrow Q$, and conversely in passing from $g: \Gamma \rightarrow Q$ to $\hat{g}: \Gamma \rightarrow G$.

3. THE SIMPLE CASE

Throughout this section, we denote by \mathbb{K} a local field, by \mathbf{G} a connected adjoint simple algebraic \mathbb{K} -group such that $G := \mathbf{G}(\mathbb{K})$ is non-compact.

Definition 3.1. Given a pair P^\pm of opposite proper parabolic subgroups of G and $z^\pm \in G/P^\pm$, a sequence g_n in G is (z^+, z^-) -contracting if g_n converges to the constant function z^+ uniformly on compact subsets of $(G/P^+) \setminus Z_{z^-}$, where $Z_{z^-} \subset G/P^+$ denotes the set of all points in G/P^+ that are opposite to z^- .

The following two lemmas are well known.

Lemma 3.2. [KLP18, Prop. 6.9] *Any unbounded sequence in G possesses a subsequence that is (z^+, z^-) -contracting for some pair P^\pm of opposite proper parabolic subgroups of G and some $z^\pm \in G/P^\pm$.*

Lemma 3.3. [GGKW17, Prop. 3.3] *Given any pair P^\pm of opposite proper parabolic subgroups of G , one can find a representation $\tau: G \rightarrow \mathrm{SL}_d(\mathbb{K})$ and τ -equivariant embeddings $\iota^\pm: G/P^\pm \rightarrow \mathrm{SL}_d(\mathbb{K})/P_1^\pm$, where P_1^+ (resp., P_1^-) denotes the stabiliser in $\mathrm{SL}_d(\mathbb{K})$ of a line (resp., a hyperplane) in \mathbb{K}^d , such that for any $z^\pm \in G/P^\pm$ and any (z^+, z^-) -contracting sequence g_n in G , the sequence $\tau(g_n)$ is $(\iota^+(z^+), \iota^-(z^-))$ -contracting.*

Lemma 3.4. *Let $h \in G$, let P^\pm be a pair of opposite proper parabolic subgroups of G , and let g_n be a (z^+, z^-) -contracting sequence in G for some $z^\pm \in G/P^\pm$. If the sequence $g_n^{-1}h g_n$ is bounded in G , then $h z^+ = z^+$.*

Proof. By Lemma 3.3, it suffices to consider the case where $G = \mathrm{SL}_d(\mathbb{K})$ and P^+ (resp., P^-) is the stabiliser in G of the line $\mathbb{K}e_1$ (resp., the hyperplane $\mathbb{K}e_2 \oplus \cdots \oplus \mathbb{K}e_d$) in \mathbb{K}^d . Let $g_n = k_n a_n k'_n$ be the KA^+K decomposition of g_n . We can assume up to subsequence that k_n (resp., k'_n) converges, say to k (resp., k'). Then a_n is a

diagonal matrix of the form $\text{diag}(\lambda_{1,n}, \dots, \lambda_{d,n})$, where $\lambda_{1,n} \geq \dots \geq \lambda_{d,n} > 0$, and since g_n is (z^+, z^-) -contracting, we have that $\lambda_{1,n}/\lambda_{2,n} \rightarrow \infty$ and $z^+ = k \cdot \mathbb{K}e_1$.

Since the sequence $g_n^{-1}hg_n = k_n^{-1}(a_n^{-1}k_n^{-1}hk_n a_n)k_n$ is bounded, so is the sequence $a_n^{-1}k_n^{-1}hk_n a_n$. Now if $(x_{1,n}, \dots, x_{d,n})^T$ is the first column of the matrix $k_n^{-1}hk_n$, then $(\frac{\lambda_{1,n}}{\lambda_{1,n}}x_{1,n}, \frac{\lambda_{1,n}}{\lambda_{2,n}}x_{2,n}, \dots, \frac{\lambda_{1,n}}{\lambda_{d,n}}x_{d,n})^T$ is the first column of the matrix $a_n^{-1}k_n^{-1}hk_n a_n$. Since $x_{i,n}$ are bounded, it follows that $x_{i,n} \rightarrow 0$ as $n \rightarrow \infty$ for $i = 2, \dots, d$. Since $k_n^{-1}hk_n \rightarrow k^{-1}hk$, it follows that $k^{-1}hk$ fixes $\mathbb{K}e_1$, and so h fixes $k \cdot \mathbb{K}e_1 = z^+$. \square

Proof of Theorem D. Suppose that D_f is not reduced to the identity and $f(\Gamma)$ is not bounded. We show that $f(\Gamma)$ is contained in a proper algebraic subgroup of G . We first note that it suffices to show that D_f is contained in a proper algebraic subgroup L of G . Indeed, by Proposition 2.5 it follows that $f(\Gamma)$ is contained in the normaliser of $\overline{\Delta_f}^{\text{Zar}}$, which is a nontrivial proper algebraic subgroup of G .

Now since $f(\Gamma)$ is not bounded in G by assumption, we have by Lemma 3.2 that there exists a pair of opposite proper parabolic subgroups $P^\pm < G$, $z^\pm \in G/P^\pm$, and a (z^+, z^-) -contracting sequence $f(\gamma_n)$ in $f(\Gamma)$. By Lemma 2.4, the sequence $f(\gamma_n)^{-1}hf(\gamma_n)$ is bounded for each $h \in D_f$. It then follows from Lemma 3.4 that $D_f \subset \text{Stab}_G(z^+)$. This completes the proof of Theorem D since $\text{Stab}_G(z^+)$ is a proper algebraic subgroup of G (indeed, a conjugate of P^+). \square

3.A. Examples. The simple case already exhibits two striking examples. Let $G = \text{PSL}_2(\mathbb{R})$, and let $F < G$ be a non-abelian free subgroup.⁵ Then F is easily seen to be Zariski-dense in G .

Example 3.5. Let $\Gamma := F$, and define a map $f: \Gamma \rightarrow G$ so that $f(g) = g$ for all $g \neq 1$ and $f(1) \neq 1$. Then $f(\Gamma)$ is unbounded and generates a group containing F , and is therefore Zariski-dense. Since f is not a homomorphism, Theorem D implies that f cannot be a quasihomomorphism.

The above example demonstrates how delicate the notion of a quasihomomorphism is; even taking a true homomorphism and changing a single value can result in a map that is not a quasihomomorphism.

Next, we exhibit a counterexample to [BV24, Theorem 4.28]⁶. The statement of the latter result is as follows. Let M be a closed negatively curved manifold, and $x_1, \dots, x_n \in \pi_1(M)$ be primitive elements that pairwise do not have conjugate powers. Let G be a non-abelian connected Lie group and let $g_1, \dots, g_n \in G$. Then it is claimed that there exists a homogeneous quasihomomorphism $\pi_1(M) \rightarrow G$ taking x_i to g_i . Recall that a quasihomomorphism is *homogeneous* if its restriction to every cyclic subgroup is a homomorphism. We exhibit a counterexample in one of the simplest cases: where M is a closed surface of genus two and $G = \text{PSL}_2(\mathbb{R})$.

Example 3.6. Let F be a discrete free subgroup of $G = \text{PSL}_2(\mathbb{R})$ of rank 4, and let $g_1, h_1, g_2, h_2 \in F$ form a free basis. Let Γ be the fundamental group of a closed surface of genus 2, and let a_1, b_1, a_2, b_2 be the standard generating set. Then the elements a_i, b_i are primitive and pairwise do not have conjugate powers, since they are independent in the abelianisation of Γ . However there is no unbounded (let alone homogeneous) quasihomomorphism $\Gamma \rightarrow G$ mapping $a_i \rightarrow g_i, b_i \rightarrow h_i, i = 1, 2$. Indeed, the image of Γ under a hypothetical such quasihomomorphism f would generate a group containing the Zariski-dense subgroup F of G , and therefore f

⁵Note that $\text{PSL}_2(\mathbb{R})$, while not itself an affine algebraic group, is the identity component in the Hausdorff topology of the \mathbb{R} -points of an adjoint simple affine algebraic group.

⁶Theorem 4.24 in the arXiv version

would necessarily be a homomorphism by Theorem D, violating the defining relation $[a_1, a_2][b_1, b_2] = 1$.

Remark 3.7. In addition to the above straightforward example, notice that unipotent groups have no non-trivial compact quotients and hence Theorem E provides a contradiction to the claim that there exist quasihomomorphisms $\Gamma \rightarrow G$ that are not constructible in the sense of Fujiwara–Kapovich when G is a non-abelian unipotent real algebraic group (cf. [BV24, Theorem 4.31]⁷).

4. REAL ALGEBRAIC GROUPS

In this section we collect some general facts about real algebraic groups that will be used in the proof of the general case. We refer the reader to [Bor91] for general background. Given a subset $X \subset G$, we denote by $Z_G(X)$ the centraliser of X , and by $N_G(X)$ the normaliser of X in G .

Lemma 4.1. *Let G be a real algebraic group. Then any nested intersection of algebraic subgroups of G eventually stabilises.*

Proof. Let $G_1 \supset G_2 \supset \dots$ be a nested intersection of algebraic subgroups of H . Eventually, the real dimension stabilises. Since a top-dimensional subgroup of a real Lie group is open, and since real algebraic groups have only finitely many connected components (see the appendix of [Mos57]), we conclude. \square

Lemma 4.2. *Let $X \subset G$ be a subset of a real algebraic group. Then there exists a finite subset $Y \subset X$ such that $Z_G(X) = Z_G(Y)$.*

Proof. Because X is a subspace of the separable metrisable space G , we have that X is itself separable. Let $\{x_1, x_2, \dots\}$ be a countable dense subset of X . Then $Z_G(X) = Z_G(\{x_1, x_2, \dots\})$. This shows that $Z_G(X)$ is the nested intersection of the algebraic subgroups $Z_G(\{x_1, \dots, x_n\})$, $n \geq 1$, so we conclude by Lemma 4.1. \square

Let $G = R \ltimes U$, where U is unipotent and R is reductive, and let $\mathfrak{g}, \mathfrak{u}, \mathfrak{r}$ denote the Lie algebras of G, U, R , respectively. We also denote by Z the centre of U and by \mathfrak{z} the Lie algebra of Z . Recall that the exponential map $\exp: \mathfrak{u} \rightarrow U$ is a homeomorphism. The following extends this slightly.

Proposition 4.3. *There exists an open neighbourhood $0 \in \omega \subset \mathfrak{g}$ such that \exp restricted to $\omega + \mathfrak{u}$ is a homeomorphism onto its image.*

The rest of this section is devoted to the proof of Proposition 4.3, and is independent from the rest of the paper.

Lemma 4.4. *For all sufficiently small $r \in \mathfrak{r}$ there exists a linear isomorphism $\varphi_r: \mathfrak{z} \rightarrow Z$ such that*

$$\exp(r + u + z) = \exp(r + u)\varphi_r(z)$$

for all $u \in \mathfrak{u}, z \in \mathfrak{z}$.

Proof. We set

$$\varphi_r(z) := \exp(-(r + u))\exp(r + u + z).$$

The main claim is that $\varphi_r(z)$ indeed depends only on r and z . Let us start by showing how this is enough to conclude. For $z_1, z_2 \in \mathfrak{z}$, we have:

$$\begin{aligned} \exp(r)\varphi_r(z_1 + z_2) &= \exp(r + z_1 + z_2) = \exp(r + z_1)\varphi_r(z_2) \\ &= \exp(r)\varphi_r(z_1)\varphi_r(z_2). \end{aligned}$$

⁷Theorem 4.27 in the arXiv version

So φ_r is additive, and since φ_r is continuous, it follows that φ_r is linear. Moreover, since \exp is a local homeomorphism, for small enough r , the map $z \mapsto \exp(r+z)$ is locally injective, hence so is φ_r , so that φ_r must therefore be a linear isomorphism.

It remains to show that $\varphi_r(z)$ is independent of u . Recall the Baker–Campbell–Hausdorff formula [Hal15, Section 5]. Let

$$B(X, Y) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \sum_{\substack{t_1+s_1>0 \\ \vdots \\ t_n+s_n>0}} \frac{[X^{t_1}Y^{s_1}X^{t_2}Y^{s_2}\dots X^{t_n}Y^{s_n}]}{\left(\sum_{j=1}^n(t_j+s_j)\right) \cdot \prod_{i=1}^n t_i!s_i!},$$

where

$$[X^{t_1}Y^{s_1}\dots X^{t_n}Y^{s_n}] = \underbrace{[X, \dots, X]}_{t_1}, \underbrace{[Y, \dots, Y]}_{s_1}, \dots, \underbrace{[X, \dots, X]}_{t_n}, \underbrace{[Y, \dots, Y]}_{s_n} \dots].$$

We set $m_n := \sum(t_j + s_j)$, which is the number of occurrences of X and Y in the above expression. The BCH formula states that

$$\exp(X)\exp(Y) = \exp(B(X, Y))$$

whenever $B(X, Y)$ converges. Note that $B(X, Y)$ indeed converges for sufficiently small X, Y .

Setting $X = -(r+u)$ and $Y = (r+u+z)$, we need to first show that the sum $B(X, Y)$ converges for all sufficiently small $r \in \mathfrak{r}$ and for all $u \in \mathfrak{u}, z \in \mathfrak{z}$. Each iterated commutator has the following form:

$$(1) \quad \pm[(r+u)^{t_1}(r+u+z)^{s_1}\dots(r+u)^{t_n}(r+u+z)^{s_n}].$$

Develop this expression using multilinearity, by separating all occurrences of r from occurrences of u and $(u+z)$. Because \mathfrak{u} is a nilpotent ideal, there is a uniform bound on the number of u or $u+z$ terms that appear in such expression, unless it vanishes. Hence we obtain a bound of the form

$$\|[(r+u)^{t_1}(r+u+z)^{s_1}\dots(r+u)^{t_n}(r+u+z)^{s_n}]\| \leq (C_{u,z}\|\mathrm{ad}(r)\|)^{m_n},$$

where $C_{u,z}$ is a constant that depends only on the nilpotency class of \mathfrak{u} and the norms $\|u\|$ and $\|u+z\|$. This proves convergence, and hence the validity of the BCH formula, for small $r \in \mathfrak{r}$ independently of u, z , as desired.

Now let us study Eq. (1) further. When $m_n = 1$, there are only two terms: $-r-u$ and $r+u+z$. They sum to z , and in particular their sum is independent of u . Therefore we need to show that when $m_n \geq 2$, the value of Eq. (1) is independent of u .

So now assume that $m_n \geq 2$. Up to grouping terms together, we may assume that $t_1, s_1 \neq 0$. Once again, we develop the expression using multilinearity, this time separating all occurrences of $(r+u)$ from occurrences of z . Up to applying the Jacobi identity, it suffices to show that

$$[r+u, \dots, [z, \dots]] = [r, \dots, [z, \dots]],$$

that is, in an iterated commutator where all terms are either $(r+u)$ or z , at least one z appears, and the first term is $(r+u)$, we may replace the first term with r . Indeed, this is because \mathfrak{z} is an ideal, and $[r+u, z'] = [r, z']$ for all $z' \in \mathfrak{z}$. \square

Proof of Proposition 4.3. We proceed by induction on the dimension of U . For the base case, if U is trivial, the statement is simply the fact that \exp is a local homeomorphism.

Now suppose that U has positive dimension, hence so does its centre Z . Let $\pi: G \rightarrow G/Z$ denote the projection. Applying the induction hypothesis on G/Z , we obtain an open neighbourhood $0 \in \omega \subset \mathfrak{g}$ such that $\exp: \mathfrak{g}/\mathfrak{z} \rightarrow G/Z$ restricted

to $d\pi(\omega) + \mathfrak{u}/\mathfrak{z}$ is a homeomorphism onto its image. We may also assume that ω is small enough that Lemma 4.4 holds for all $r \in \mathfrak{t} \cap \omega$. We are left to prove that $\exp: \omega + \mathfrak{u} \rightarrow \exp(\omega)U$ is a homeomorphism.

Let $v_1, v_2 \in \omega + \mathfrak{u}$ be such that $\exp(v_1) = \exp(v_2)$. Then $d\pi(v_1) = d\pi(v_2)$, so there exists $z \in \mathfrak{z}$ such that $v_2 = v_1 + z$. We write $v_1 = r + u$, for $r \in \mathfrak{t} \cap \omega$. Hence $v_2 = r + u + z$, and by Lemma 4.4 we have $\exp(v_2) = \exp(v_1)\varphi_r(z)$. Because φ_r is injective, $z = 0$, that is $v_1 = v_2$.

Let $g \in \exp(\omega)U$. By induction there exists $v = r + u \in (\mathfrak{t} \cap \omega) + \mathfrak{u}$ such that $\pi(g) = \exp(d\pi(v))$. So there exists $z \in Z$ such that $g = \exp(v)z$. Choose $z' \in \mathfrak{z}$ such that $\varphi_r(z') = z$. Then

$$g = \exp(r + u)\varphi_r(z) = \exp(r + u + z) \in \exp(\omega + \mathfrak{u}). \quad \square$$

5. PROOF OF THE MAIN THEOREM

In this section we prove the main theorem in its various versions (Theorem A, Theorem B, Corollary C and Theorem E). Let $f: \Gamma \rightarrow G$ be a quasihomomorphism, where G is a real algebraic group. Let $D = D_f$ denote the defect set, and $\Delta = \Delta_f$ the defect group, i.e. the closure of the group generated by D .

5.A. Setup.

Lemma 5.1. *Up to bounded modification, G is Zariski-connected, and $\langle f(\Gamma) \rangle$ is Zariski-dense.*

This will be our only bounded modification, until Claim 5.8. We also remark that this is the only step where we need to pass to a finite-index subgroup of Γ .

Proof. Replacing G by the Zariski-closure of $\langle f(\Gamma) \rangle$ is clearly a bounded modification. Next, we will show that there exists a bounded modification that takes values in the Zariski-connected component of G . By Lemma 4.1, we can repeat this process until both properties hold at once.

Let $1 \in G_0, G_1, \dots, G_n$ denote the Zariski-connected components of G . Let $X_i = f^{-1}(G_i \cap f(\Gamma)) \subset \Gamma$. By [BHS19, Lemma 7]⁸, there exists $i \in \{0, \dots, n\}$ and $m \geq 1$ such that $(X_i X_i^{-1})^m$ is a finite-index subgroup Γ' of Γ . Now, for an element $x \in \Gamma'$ choose an expression $x = y_1 z_1^{-1} \cdots y_m z_m^{-1}$, where $y_j, z_j \in X_i$. Define

$$f'(x) = f(y_1)f(z_1)^{-1} \cdots f(y_m)f(z_m)^{-1}$$

and notice that $f(y_j)f(z_j)^{-1} \in G_i G_i^{-1} \subset G_0$, so $f'(\Gamma') \subset G_0$. That f' is a quasihomomorphism close to $f|_{\Gamma'}$ follows from Lemma 2.6. Therefore f' is a bounded modification of f that takes values in G_0 . \square

Now G is a Zariski-connected real algebraic group, so it decomposes as $G = (R/F) \times U$, where U is unipotent, R is reductive, and F is a finite subgroup of R . The action of R/F on U pulls back to an action of R , which defines a semidirect product $R \ltimes U$, which surjects on H with kernel F . By Remark 2.7, we can therefore assume that

$$G = R \times U \quad \text{with } R \text{ connected reductive and } U \text{ unipotent.}$$

We now decompose the reductive part as

$$R = G_b \times G_h \times T,$$

where

- T is a torus;

⁸The cited lemma is for locally compact second countable groups, but this hypothesis is only used for the compactness of the Chabauty topology on the set of closed subsets, which holds in the discrete case without any cardinality assumptions.

- G_b is the product of all those simple factors S such that the composition $\Gamma \xrightarrow{f} G \rightarrow S$ is bounded but does not have central defect;
- G_h is the product of all those simple factors S such that the composition $\Gamma \xrightarrow{f} G \rightarrow S$ has central defect.

By Theorem D, every simple factor is indeed either in G_b or in G_h . We also note that the projection of $\langle D \rangle$ to G_h is trivial, hence so is the projection of Δ to G_h .

5.B. **The conjugacy action on Δ .** Recall from Lemma 2.4 that for all $x \in \langle D \rangle$, the orbit $x^{f(\Gamma)}$ is bounded. It is a priori not clear that this extends to $x \in \Delta$, or to $x \in \overline{\Delta}^{\text{Zar}}$. Note that this is a feature unique to our setting and does not occur when the target is discrete, as in [FK16]. Our next goal is to tackle Δ .

Lemma 5.2. *For every identity neighbourhood $W \subset \Delta$, there exists an identity neighbourhood $W_0 \subset W$ such that $W_0^{f(\Gamma)} \subset W$. Moreover, for every $x \in \Delta$, the set $x^{f(\Gamma)}$ is bounded.*

Again, the “moreover” part holds for $x \in \langle D \rangle$, by Lemma 2.4. Indeed, choosing W to be a compact identity neighbourhood of Δ , Lemma 5.2 gives the same result for the corresponding W_0 . So $x^{f(\Gamma)}$ is bounded for all $x \in W_0 \langle D \rangle = \Delta$, as claimed. Hence we can focus on the first statement.

Using the same notation as in Proposition 4.3, we fix an open neighbourhood $0 \in \omega \subset \mathfrak{g}$ such that \exp restricted to $\omega + \mathfrak{u}$ is a homeomorphism onto its image.

Claim 5.3. *There exists an open neighbourhood $0 \in \omega' \subset \mathfrak{g}$ such that*

$$\text{Ad}(f(\Gamma))(\exp^{-1}(\Delta) \cap \omega' + \mathfrak{u}) \subset \omega + \mathfrak{u}.$$

Proof. Throughout the proof, we work modulo \mathfrak{u} , hence in the Lie algebra \mathfrak{r} of R . Recall that R splits as $G_b \times G_h \times T$, that the projection of $f(\Gamma)$ onto G_b is bounded, the projection of Δ onto G_h is finite and central, and T is abelian. The Lie algebra \mathfrak{r} has an Ad-invariant splitting as the direct sum of the Lie algebras of the three factors, and we study the adjoint action coordinate-wise.

Because T is central in R , we have that T is in the kernel of the adjoint action. Because Δ has finite central projection onto G_h , the preimage $\exp^{-1}(\Delta)$ has trivial \mathfrak{g}_h component: indeed, non-trivial central elements of the semi-simple group G_h cannot be in the image of the exponential map. So the adjoint action of $f(\Gamma)$ on $\exp^{-1}(\Delta)$ reduces to the adjoint action of the projection of $f(\Gamma)$ onto G_b , which is bounded by definition. Hence $\text{Ad}(f(\Gamma))$ is a uniformly bounded family of operators of $\exp^{-1}(\Delta)$. It follows that the desired open neighbourhood ω' of 0 exists. \square

Define

$$\mathfrak{b} := \{v \in \mathfrak{g} : \text{Ad}(f(\Gamma))(v) \text{ is bounded}\}.$$

Claim 5.4. $\Delta \cap \exp(\omega' + \mathfrak{u}) \subset \exp(\mathfrak{b})$.

Proof. Suppose first that $x \in \langle D \rangle \cap \exp(\omega' + \mathfrak{u})$. Let $v \in \omega' + \mathfrak{u}$ be such that $\exp(v) = x$. Then

$$x^{f(\Gamma)} = \exp(\text{Ad}(f(\Gamma))(v)).$$

By Claim 5.3:

$$\text{Ad}(f(\Gamma))(v) \subset \omega + \mathfrak{u}.$$

Because $\exp|_{\omega + \mathfrak{u}}$ is a homeomorphism onto its image, and $x^{f(\Gamma)}$ is bounded by Lemma 2.4, $\text{Ad}(f(\Gamma))(v)$ must be bounded as well, that is $v \in \mathfrak{b}$.

Now for the general case, suppose that $x \in \Delta \cap \exp(\omega' + \mathfrak{u})$, and let $x_n \in \langle D \rangle$ be such that $x_n \rightarrow x$. Up to discarding finitely many terms, we may assume that $x_n \in \exp(\omega' + \mathfrak{u})$, so by the previous case we can choose $v_n \in (\omega' + \mathfrak{u}) \cap \mathfrak{b}$ such

that $\exp(v_n) = x_n$. Because \exp is a homeomorphism on $\omega' + \mathfrak{u}$, we have that v_n converges to some $v \in \omega' + \mathfrak{u}$ such that $\exp(v) = x$. Finally, since \mathfrak{b} is a vector subspace of \mathfrak{g} and is in particular closed, we have that $v_n \in \mathfrak{b}$ implies $v \in \mathfrak{b}$. \square

Proof of Lemma 5.2. Let \mathfrak{d} denote the Lie algebra of Δ . Claim 5.4 implies that $\mathfrak{d} \subset \mathfrak{b}$. Let W be an identity neighbourhood of Δ , so $\exp^{-1}(W)$ is an open subset of \mathfrak{d} . Because $\text{Ad}(f(\Gamma))$ is a bounded family of operators of \mathfrak{b} , hence of \mathfrak{d} , there exists an open subset ω_0 of \mathfrak{d} such that $\text{Ad}(f(\Gamma))(\omega_0) \subset \exp^{-1}(W)$. Taking $W_0 = \exp(\omega_0)$, we have

$$W_0^{f(\Gamma)} = \exp(\text{Ad}(f(\Gamma))(\omega_0)) \subset W. \quad \square$$

5.C. The conjugacy action on $\overline{\Delta}^{\text{Zar}}$. In this subsection we strengthen and extend Lemma 5.2 to apply to the Zariski-closure $\overline{\Delta}^{\text{Zar}}$.

Proposition 5.5. *There exists a compact set $K \subset G$ such that $f(\Gamma) \subset Z_G(\overline{\Delta}^{\text{Zar}})K$. In particular, for all $x \in \overline{\Delta}^{\text{Zar}}$, the set $x^{f(\Gamma)}$ is bounded.*

By definition $\Delta = \langle \overline{D} \rangle$ is the closure of a group generated by a bounded set and is hence compactly generated. So we can apply [Hoc52] to see that the group of automorphisms $\text{Aut}(\Delta)$ with the compact open topology is a Lie group. A set of the form

$$\mathcal{N}(C, W) := \{\varphi \in \text{Aut}(\Delta) : \varphi(x) \in xW \text{ for all } x \in C\},$$

for C a compact set and W an open identity neighbourhood, is an identity neighbourhood.

We now consider the map

$$\alpha: N_G(\Delta) \rightarrow \text{Aut}(\Delta)$$

given by the conjugacy action.

Claim 5.6. *The set $\alpha(f(\Gamma))$ is bounded in $\text{Aut}(\Delta)$.*

Proof. We verify this using the Arzela–Ascoli Theorem. First, for every $x \in \Delta$, the second part of Lemma 5.2 gives that $\alpha(f(\Gamma))(x) = x^{f(\Gamma)}$ is bounded. Moreover, $\alpha(f(\Gamma))$ is equicontinuous: because this is a group topology, it suffices to verify this locally, which is given by the first part of Lemma 5.2. \square

Claim 5.7. *There exists a finite subset $F \subset \Delta$, an open identity neighbourhood $W \subset \Delta$, and a compact set $K \subset G$ such that*

$$\alpha^{-1}(\mathcal{N}(F, W)) \subset Z_G(\overline{\Delta}^{\text{Zar}})K.$$

Proof. Because centralising is an algebraic condition, $Z_G(\Delta) = Z_G(\overline{\Delta}^{\text{Zar}})$. By Lemma 4.2 we can choose $F = \{x_1, \dots, x_n\} \subset \Delta$ to be such that $Z_G(F) = Z_G(\Delta)$.

Consider the action of G on the product G^n by coordinate-wise conjugacy, and let O denote the orbit of (x_1, \dots, x_n) . By [Zim84, Theorem 3.1.3], the action has locally closed orbits. In particular, there exists a compact neighbourhood W of e in G such that

$$O \cap (x_1W, \dots, x_nW)$$

is compact in O . Moreover, by [Zim84, Theorem 2.1.14] the orbit map

$$G/Z_G(F) \rightarrow O$$

is a homeomorphism. Therefore the preimage of $O \cap (x_1W, \dots, x_nW)$ is compact in $G/Z_G(F)$, and we can write its lift to G as $Z_G(F)K$, for a compact subset $K \subset G$.

Now if $g \in \alpha^{-1}(\mathcal{N}(F, W))$, then for all $i = 1, \dots, n$, we have

$$x_i^g = \alpha(g)(x_i) \in x_iW,$$

that is g lies in the preimage of $O \cap (x_1W, \dots, x_nW)$ under the orbit map, and so it belongs to $Z_G(F)K = Z_G(\overline{\Delta}^{\text{Zar}})K$ by the above. \square

Proof of Proposition 5.5. Let F, W be as in Claim 5.7. By Claim 5.6, there exist finitely many elements $\gamma_1, \dots, \gamma_n \in \Gamma$ such that

$$\alpha(f(\Gamma)) \subset \bigcup_i \mathcal{N}(F, W)\alpha(f(\gamma_i)),$$

hence

$$f(\Gamma) \subset \bigcup_i \alpha^{-1}(\mathcal{N}(F, W))f(\gamma_i).$$

By Claim 5.7, we therefore have

$$f(\Gamma) \subset Z_H(\overline{\Delta}^{\text{Zar}})K\{f(\gamma_1), \dots, f(\gamma_n)\}.$$

Since $K\{f(\gamma_1), \dots, f(\gamma_n)\}$ is still compact, we conclude. \square

5.D. Endgame. By Proposition 5.5, there exists a compact set K such that $f(\Gamma) \subset Z_G(\overline{\Delta}^{\text{Zar}})K$. Define $N := \overline{\Delta}^{\text{Zar}}Z_G(\overline{\Delta}^{\text{Zar}})$, which is an algebraic normal subgroup of G containing the defect. Therefore the composition

$$\Gamma \xrightarrow{f} G \rightarrow G/N$$

is a homomorphism. Its image is a precompact Zariski-dense subgroup of the algebraic group G/N , which implies that G/N itself is compact⁹ In particular N must contain the unipotent U , and so we can write

$$G = BN,$$

for B a compact algebraic subgroup of G contained in R .

Now for each $\gamma \in \Gamma$ we can choose a decomposition of the form

$$f(\gamma) = b(\gamma)z(\gamma)d(\gamma) \in B \cdot Z_G(\overline{\Delta}^{\text{Zar}}) \cdot \overline{\Delta}^{\text{Zar}}.$$

Because $f(\Gamma) \subset Z_G(\overline{\Delta}^{\text{Zar}})K$, and $Z_G(\overline{\Delta}^{\text{Zar}})$ is normal, we may choose this expression so that $d(\Gamma)$ is bounded. Set $f'(\gamma) := b(\gamma)z(\gamma) \in BZ_G(\overline{\Delta}^{\text{Zar}})$.

Claim 5.8. $f' : \Gamma \rightarrow BZ_G(\overline{\Delta}^{\text{Zar}})$ is a bounded modification of f . Moreover, $D_{f'} \subset \overline{\Delta}^{\text{Zar}}$.

Proof. We have that f' is close to f because $d(\Gamma)$ is bounded. Recall from Notation 2.1, that for a bounded set K , we write $x \approx_K y$ if $x \in yK$. In the course of the proof, we write \approx_f if $K = D_f$, \approx_d if $K = d(\Gamma)$, and finally $\approx_{B,d}$ if $K = \{bdb^{-1} : b \in B, d \in d(\Gamma)\}$. Note that each of these bounded sets is contained in $\overline{\Delta}^{\text{Zar}}$.

Now we estimate:

$$\begin{aligned} f'(\gamma\delta) &\approx_d f(\gamma\delta) \approx_f f(\gamma)f(\delta) \approx_d f(\gamma)f'(\delta) \\ &= f'(\gamma)d(\gamma)b(\delta)z(\delta) = f'(\gamma)b(\delta)d(\gamma)^{b(\delta)}z(\delta) \\ &= f'(\gamma)b(\delta)z(\delta)d(\gamma)^{b(\delta)} \approx_{B,d} f'(\gamma)f'(\delta). \end{aligned}$$

Here we used that $\overline{\Delta}^{\text{Zar}}$ is normal; hence $d(\gamma)^{b(\delta)} \subset \overline{\Delta}^{\text{Zar}}$ commutes with $z(\delta)$.

Finally, as we remarked above, each compact set appearing in an approximate equality \approx is contained in $\overline{\Delta}^{\text{Zar}}$, so $D_{f'} \subset \overline{\Delta}^{\text{Zar}}$. \square

⁹This is the only step where we use the hypothesis that G is a *real* algebraic group, as opposed to a *p-adic* algebraic group.

Proof of Theorem A. By Claim 5.8, we may consider

$$f' : \Gamma \rightarrow BZ_G(\overline{\Delta}^{\text{Zar}}) = Z_G(\overline{\Delta}^{\text{Zar}})B,$$

where B is a compact (algebraic) subgroup, and $D' := D_{f'}$ is contained in $\overline{\Delta}^{\text{Zar}}$. Therefore

$$(D')^G = (D')^{Z_G(\overline{\Delta}^{\text{Zar}})B} = (D')^B \subset \overline{\Delta}^{\text{Zar}}.$$

So $(D')^B$ is a normal bounded subset of G containing the defect D' . \square

Proof of Theorem E. Suppose that $\langle f(\Gamma) \rangle$ is Zariski-dense in G , and G has no non-trivial compact quotient. Then G is automatically Zariski-connected, therefore the bounded modification from Lemma 5.1 is not needed in this case. Now with the notation above, we have that B is a compact quotient of G , so B must be trivial, which implies that the bounded modification from Claim 5.8 takes values in the centraliser of the defect. \square

Proof of Theorem B. As in the proof of Theorem A, we have a bounded modification $f' : \Gamma \rightarrow BZ_G(\overline{\Delta}^{\text{Zar}})$, where B is a compact (algebraic) subgroup, and $D_{f'}$ is contained in $\overline{\Delta}^{\text{Zar}}$. The action of G' by conjugacy on $\overline{\Delta}^{\text{Zar}}$ factors through B , hence the action of $\overline{\Delta}^{\text{Zar}}$ by conjugacy on itself factors through a compact group C_0 . This implies that

$$\overline{\Delta}^{\text{Zar}} = Z_\Delta \times C_0,$$

where Z_Δ is the centre of $\overline{\Delta}^{\text{Zar}}$, and C_0 is a compact group. It follows that there exists a unique maximal compact (algebraic) subgroup $C < \overline{\Delta}^{\text{Zar}}$, which is characteristic in $\overline{\Delta}^{\text{Zar}}$ hence normal in G' . Quotienting by C , the image of $\overline{\Delta}^{\text{Zar}}$ is a simply connected abelian normal subgroup $A \triangleleft G'/C$, and the action by conjugacy of G' on A factors through the compact group B , so that A is rigid in G'/C . \square

Proof of Corollary C. Let us start by noticing that if $H_b^2(\Gamma; \pi)$ vanishes for every finite-dimensional orthogonal representation π with no non-trivial invariant vectors, and vanishes when π is the trivial representation, then $H_b^2(\Gamma; \pi)$ vanishes for every finite-dimensional orthogonal representation π [Mon01, Lemma 1.2.10].

Thanks to Theorem B and Remark 2.7, we may reduce to the case in which $f : \Gamma \rightarrow G$ is such that D_f is contained in a rigid abelian algebraic normal subgroup $A \triangleleft G$, such that $D = D_f$ is contained in A . We will show that, under the assumption that bounded cohomology vanishes with coefficients in every finite-dimensional orthogonal representation, there exists a bounded map $b : \Gamma \rightarrow A$ such that $\gamma \mapsto f(\gamma)b(\gamma)$ is a homomorphism. This will conclude the proof of the second statement, since the map $\gamma \mapsto f(\gamma)b(\gamma)$ is a bounded modification of f .

The composition $\Gamma \xrightarrow{f} G \rightarrow G/A$ is a homomorphism, therefore defines an orthogonal representation π of Γ on A . Now consider the lifting problem

$$\begin{array}{ccc} & & G \\ & \nearrow \text{---} & \downarrow \\ \Gamma & & G/A \\ & \searrow & \end{array}$$

The set-theoretic lift $f : \Gamma \rightarrow G$ defines the bounded 2-cocycle

$$(\gamma_1, \gamma_2) \mapsto f(\gamma_2)^{-1}f(\gamma_1)^{-1}f(\gamma_1\gamma_2) \in A,$$

which in turn represents a class in $H_b^2(\Gamma; \pi)$. The stronger vanishing hypothesis gives a bounded primitive $b: \Gamma \rightarrow A$ for this cocycle, which means exactly that $\gamma \mapsto f(\gamma)b(\gamma)$ is a homomorphism.

Suppose instead that we only have vanishing for representations with no non-trivial invariant vectors. Then $H_b^i(\Gamma; (\pi^\Gamma)^\perp)$ vanishes for $i = 1$ (this is always true [Mon01, Proposition 6.2.1]) and $i = 2$, so the long exact sequence in bounded cohomology gives an isomorphism $H_b^2(\Gamma; \pi^\Gamma) \rightarrow H_b^2(\Gamma; \pi)$ induced by the change of coefficients map. That is, every bounded 2-cocycle is boundedly cohomologous to one that takes values in the invariants. Therefore the same argument as above shows that there exists a bounded map $b: \Gamma \rightarrow A$ such that $\gamma \mapsto f(\gamma)b(\gamma)$ has central defect. \square

6. COUNTEREXAMPLE TO CENTRAL DEFECT

Proposition 6.1. *Let Γ be an acylindrically hyperbolic group that surjects onto \mathbb{Z} . Let $G = \mathrm{SO}(2) \times \mathbb{R}^2$. Then there exists a quasihomomorphism $f: \Gamma \rightarrow G$ such that no bounded modification of f has central defect.*

Acylindrically hyperbolic groups include non-abelian free groups, and more generally all non-elementary (relatively) hyperbolic groups [Osi18].

We use again the exponential notation for conjugacy; note that for $v \in \mathbb{R}^2$ and $A \in \mathrm{SO}(2)$, the conjugacy in G written as v^A is just the ordinary right action of A on v .

Proof. The map to \mathbb{Z} induces a homomorphism $u: \Gamma \rightarrow \mathrm{SO}(2)$ with dense image. Recall that a u -quasicocycle is a map $c: \Gamma \rightarrow \mathbb{R}^2$ such that

$$D_c := \{c(gh) - (c(g))^{u(h)} + c(h) : g, h \in \Gamma\}$$

is bounded. If D_c is trivial, we call c a u -cocycle. We can choose a u -quasicocycle c that is not close to any u -cocycle. By [BBF16, Theorem 1.1], this is possible provided that there exists an invariant vector for the maximal finite normal subgroup of Γ ; in our case, we chose an action that factors through \mathbb{Z} , hence every finite subgroup of Γ in fact acts trivially.

Define $f = (u, c): \Gamma \rightarrow \mathrm{SO}(2) \times \mathbb{R}^2$. This is a quasihomomorphism:

$$\begin{aligned} f(gh) &= (u(gh), c(gh)) \approx_{D_c} (u(g)u(h), c(g)^{u(h)} + c(h)) \\ &= (u(g), c(g))(u(h), c(h)) = f(g)f(h). \end{aligned}$$

Suppose by contradiction that there exists a bounded modification $f': \Gamma' \rightarrow G'$ such that G' centralises $D_{f'}$. Note that $u|_{\Gamma'}: \Gamma' \rightarrow \mathrm{SO}(2)$ is still a homomorphism with dense image, and $c|_{\Gamma'}$ is a $u|_{\Gamma'}$ quasicocycle. Moreover, it follows from the interpretation of quasicocycles in terms of bounded cohomology, and the injectivity of the restriction to Γ' [Mon01, Proposition 8.6.2], that $c|_{\Gamma'}$ is not close to any $u|_{\Gamma'}$ -cocycle. Therefore, we may assume that $\Gamma' = \Gamma$, and for clarity we will denote $f' = \varphi$, and $G' = H$. By assumption, there exists a bounded set $D_{\varphi, f}$ such that $\varphi(g) \approx_{D_{\varphi, f}} f(g)$ (see Notation 2.1).

Suppose first that H abelian. Then φ is a quasimorphism with values in an abelian group, and in particular φ is uniformly bounded on commutators, more precisely $\varphi([g, h])$ is bounded by twice the defect D_φ [Cal09, Section 2.2.3].

$$\begin{aligned} 0 &\approx_{2D_\varphi} \varphi([g, h]) \approx_{D_{\varphi, f}} f([g, h]) = c([g, h]) = c(g^{-1}h^{-1}gh) \\ &\approx_{D_c} c(g^{-1})^{u(h^{-1}gh)} + c(h^{-1})^{u(gh)} + c(g)^{u(h)} + c(h). \end{aligned}$$

Now

$$\begin{aligned} 0 &\approx_{D_f} f(gg^{-1}) = c(gg^{-1}) \approx_{D_c} c(g)^{u(g^{-1})} + c(g^{-1}) \\ &\Rightarrow c(g^{-1}) \approx_{D_c \cdot D_f} -c(g)^{u(g^{-1})}. \end{aligned}$$

Combining the above, together with the fact that u is a homomorphism to an abelian group, we get a bounded set K such that

$$\begin{aligned} 0 &\approx_K -c(g) - c(h)^{u(g)} + c(g)^{u(h)} + c(h) \\ &\Rightarrow c(g)^{I-u(h)} \approx_K c(h)^{I-u(g)}. \end{aligned}$$

Fixing g and letting h vary, the left hand side remains bounded in terms of g , which implies that $c(\Gamma)$ is uniformly close to the fixed point set of $u(g)$, which is $\{0\}$ as soon as $u(g)$ is non-trivial. Therefore $c(\Gamma)$ is bounded, contradiction.

Now suppose that H is not abelian. Then H cannot be contained in \mathbb{R}^2 , so there exists $(x, a) \in H$ with $x \neq I$. The centraliser of (x, a) intersects \mathbb{R}^2 trivially, which implies that D_φ intersects \mathbb{R}^2 trivially. Similarly, H cannot project injectively to $\text{SO}(2)$, so there exists $(I, b) \in H$ with $b \neq 0$. The centraliser of (I, b) is contained in \mathbb{R}^2 , and therefore $D_\varphi \subset \mathbb{R}^2$. We conclude that D_φ is trivial, i.e. that φ is a homomorphism. So we can write $\varphi(g) = (v(g), d(g))$, where $v: \Gamma \rightarrow \text{SO}(2)$ is a homomorphism, and $d: \Gamma \rightarrow \mathbb{R}^2$ is a v -cocycle, i.e. $d(gh) = d(g)^{v(h)} + d(h)$. If $v = u$, then d will be a u -cocycle, and $\varphi(g) \approx_{D_{\varphi, f}} f(g) \Rightarrow d(g) \approx_{D_{\varphi, f}} c(g)$, which contradicts our choice of c . Therefore, it remains to show that $v = u$.

Write $\tilde{d}(g) = d(g)^{v(g)^{-1}}$. Then

$$\tilde{d}(gh) = d(gh)^{v(gh)^{-1}} = (d(g)^{v(h)} + d(h))^{v(gh)^{-1}} = \tilde{d}(g) + \tilde{d}(h)^{v(g)^{-1}}.$$

In words, \tilde{d} is a *left* v -cocycle, where we consider the left action of $\text{SO}(2)$ on \mathbb{R}^2 given by right multiplication with the inverse. Moreover,

$$\varphi(g) = (v(g), d(g)) = \tilde{d}(g)v(g) \approx_{\text{SO}(2)} \tilde{d}(g).$$

Similarly, $\tilde{c}(g) = c(g)^{u(g)^{-1}}$ is a left u -quasicocycle, and $f(g) \approx_{\text{SO}(2)} \tilde{c}(g)$. Therefore $\tilde{c}(g) \approx_{\text{SO}(2)} \tilde{d}(g)$. We now estimate:

$$\begin{aligned} \tilde{c}(g) + \tilde{c}(h)^{u(g)^{-1}} &\approx_{D_{\tilde{c}}} \tilde{c}(gh) \approx_{\text{SO}(2)} \tilde{d}(gh) \\ &= \tilde{d}(g) + \tilde{d}(h)^{v(g)^{-1}} \approx_{\text{SO}(2)} \tilde{c}(g) + \tilde{c}(h)^{v(g)^{-1}}. \end{aligned}$$

We have shown that there exists a bounded set K such that

$$\tilde{c}(h)^{u(g)^{-1}} \approx_K \tilde{c}(h)^{v(g)^{-1}}.$$

Note that $\tilde{c}(\Gamma)$ is unbounded, because $f(\Gamma)$ is unbounded. Fixing g and letting h vary, we conclude that $v = u$, which finishes the proof. \square

Of course, Proposition 6.1 does not contradict Theorem A, because the defect of f is contained in \mathbb{R}^2 , and so its $\text{SO}(2)$ -closure is bounded and normal.

REFERENCES

- [BBF16] Mladen Bestvina, Ken Bromberg, and Koji Fujiwara. Bounded cohomology with coefficients in uniformly convex Banach spaces. *Comment. Math. Helv.*, 91(2):203–218, 2016. [10.4171/CMH/383](#). Cited on Page 14.
- [BF02] Mladen Bestvina and Koji Fujiwara. Bounded cohomology of subgroups of mapping class groups. *Geom. Topol.*, 6:69–89, 2002. [10.2140/gt.2002.6.69](#). Cited on Page 1.
- [BHS19] Michael Björklund, Tobias Hartnick, and Thierry Stulemeijer. Borel density for approximate lattices. *Forum Math. Sigma*, 7:Paper No. e40, 27, 2019. [10.1017/fms.2019.39](#). Cited on Page 9.
- [BM99] M. Burger and N. Monod. Bounded cohomology of lattices in higher rank Lie groups. *J. Eur. Math. Soc. (JEMS)*, 1(2):199–235, 1999. [10.1007/s100970050007](#). Cited on Page 1.

- [BM02] M. Burger and N. Monod. Continuous bounded cohomology and applications to rigidity theory. *Geom. Funct. Anal.*, 12(2):219–280, 2002. [10.1007/s00039-002-8245-9](https://doi.org/10.1007/s00039-002-8245-9). Cited on Page 1.
- [Bog25] Elena Bogliolo. Bounded cohomology and scl of verbal wreath products, 2025. [arXiv:2501.07438](https://arxiv.org/abs/2501.07438) [[math.GR](#)]. Cited on Page 3.
- [Bor91] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991. [10.1007/978-1-4612-0941-6](https://doi.org/10.1007/978-1-4612-0941-6). Cited on Page 7.
- [BV24] Michael Brandenbursky and Misha Verbitsky. Non-commutative barge-ghys quasi-morphisms. *IMRN*, may 2024. Ahead of press, [10.1093/imrn/rnae119](https://doi.org/10.1093/imrn/rnae119). Cited on Pages 2, 6, and 7.
- [Cal09] Danny Calegari. *scl*, volume 20 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2009. [10.1142/e018](https://doi.org/10.1142/e018). Cited on Pages 1 and 14.
- [CFFLM25] Caterina Campagnolo, Francesco Fournier-Facio, Yash Lodha, and Marco Moraschini. An algebraic criterion for the vanishing of bounded cohomology, 2025. [arXiv:2311.16259](https://arxiv.org/abs/2311.16259) [[math.GR](#)], to appear in *Compos. Math.* Cited on Page 3.
- [Dum24] Guillaume Dumas. On quasi-homomorphism rigidity for lattices in simple algebraic groups, 2024. [arXiv:2403.16500](https://arxiv.org/abs/2403.16500) [[math.FA](#)]. Cited on Page 1.
- [FK16] K. Fujiwara and M. Kapovich. On quasihomomorphisms with noncommutative targets. *Geom. Funct. Anal.*, 26(2):478–519, 2016. [10.1007/s00039-016-0364-9](https://doi.org/10.1007/s00039-016-0364-9). Cited on Pages 1, 2, 3, 4, and 10.
- [Fri17] R. Frigerio. *Bounded cohomology of discrete groups*, volume 227 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2017. [10.1090/surv/227](https://doi.org/10.1090/surv/227). Cited on Page 1.
- [GGKW17] François Guéritaud, Olivier Guichard, Fanny Kassel, and Anna Wienhard. Anosov representations and proper actions. *Geom. Topol.*, 21(1):485–584, 2017. [10.2140/gt.2017.21.485](https://doi.org/10.2140/gt.2017.21.485). Cited on Page 5.
- [Ghy87] E. Ghys. Groupes d’homéomorphismes du cercle et cohomologie bornée. In *The Lefschetz centennial conference, Part III (Mexico City, 1984)*, volume 58 of *Contemp. Math.*, pages 81–106. Amer. Math. Soc., Providence, RI, 1987. [10.24033/msmf.358](https://doi.org/10.24033/msmf.358). Cited on Page 1.
- [Hal15] Brian Hall. *Lie groups, Lie algebras, and representations: An elementary introduction*, volume 222 of *Graduate Texts in Mathematics*. Springer, Cham, second edition, 2015. [10.1007/978-3-319-13467-3](https://doi.org/10.1007/978-3-319-13467-3). Cited on Page 8.
- [Hoc52] Gerhard Hochschild. The automorphism group of a Lie group. *Transactions of the American Mathematical Society*, 72(2):209–216, 1952. [10.1090/S0002-9947-1952-0045735-2](https://doi.org/10.1090/S0002-9947-1952-0045735-2). Cited on Page 11.
- [Hru22] Ehud Hrushovski. Beyond the Lascar Group, 2022. [arXiv:2011.12009](https://arxiv.org/abs/2011.12009) [[math.LO](#)]. Cited on Pages 1 and 2.
- [Joh72] Barry Edward Johnson. *Cohomology in Banach algebras*. Memoirs of the American Mathematical Society, No. 127. American Mathematical Society, Providence, RI, 1972. Cited on Page 3.
- [KLP18] Michael Kapovich, Bernhard Leeb, and Joan Porti. Dynamics on flag manifolds: domains of proper discontinuity and cocompactness. *Geom. Topol.*, 22(1):157–234, 2018. [10.2140/gt.2018.22.157](https://doi.org/10.2140/gt.2018.22.157). Cited on Page 5.
- [Mal04] A. V. Maljutin. Writhe of (closed) braids. *Algebra i Analiz*, 16(5):59–91, 2004. [10.1090/S1061-0022-05-00879-4](https://doi.org/10.1090/S1061-0022-05-00879-4). Cited on Page 1.
- [Man08] Jason Fox Manning. Actions of certain arithmetic groups on Gromov hyperbolic spaces. *Algebr. Geom. Topol.*, 8(3):1371–1402, 2008. [10.2140/agt.2008.8.1371](https://doi.org/10.2140/agt.2008.8.1371). Cited on Page 1.
- [Mon01] Nicolas Monod. *Continuous bounded cohomology of locally compact groups*, volume 1758 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2001. [10.1007/b80626](https://doi.org/10.1007/b80626). Cited on Pages 13 and 14.
- [Mon22] Nicolas Monod. Lamplighters and the bounded cohomology of Thompson’s group. *Geom. Funct. Anal.*, 32(3):662–675, 2022. [10.1007/s00039-022-00604-9](https://doi.org/10.1007/s00039-022-00604-9). Cited on Page 3.
- [Mos57] G. D. Mostow. On the fundamental group of a homogeneous space. *Ann. of Math. (2)*, 66:249–255, 1957. [10.2307/1969997](https://doi.org/10.2307/1969997). Cited on Page 7.
- [MS04] Nicolas Monod and Yehuda Shalom. Cocycle superrigidity and bounded cohomology for negatively curved spaces. *J. Differential Geom.*, 67(3):395–455, 2004. [10.4310/jdg/1102091355](https://doi.org/10.4310/jdg/1102091355). Cited on Page 3.

- [Osi18] Denis V. Osin. Groups acting acylindrically on hyperbolic spaces. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures*, pages 919–939. World Sci. Publ., Hackensack, NJ, 2018. Cited on Page 14.
- [Oza11] Narutaka Ozawa. Quasi-homomorphism rigidity with non-commutative targets. *J. Reine Angew. Math.*, 655:89–104, 2011. [10.1515/CRELLE.2011.034](https://doi.org/10.1515/CRELLE.2011.034). Cited on Page 1.
- [PR14] L. Polterovich and D. Rosen. *Function theory on symplectic manifolds*, volume 34 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 2014. [10.1090/crmm/034](https://doi.org/10.1090/crmm/034). Cited on Page 1.
- [Tao14] Terence Tao. *Hilbert’s fifth problem and related topics*, volume 153 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2014. [10.1090/gsm/153](https://doi.org/10.1090/gsm/153). Cited on Page 3.
- [Zim84] R. J. Zimmer. *Ergodic theory and semisimple groups*, volume 81 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1984. [10.1007/978-1-4684-9488-4](https://doi.org/10.1007/978-1-4684-9488-4). Cited on Page 11.

(S. Douba) MATHEMATISCHES INSTITUT, RHEINISCHE FRIEDRICH-WILHELMS-UNIVERSITÄT BONN,
ENDENICHER ALLEE 60, 53115 BONN, GERMANY
Email address: douba@math.uni-bonn.de

(F. Fournier-Facio) DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS,
UNIVERSITY OF CAMBRIDGE, UK
Email address: ff373@cam.ac.uk

(S. Hughes) MATHEMATISCHES INSTITUT, RHEINISCHE FRIEDRICH-WILHELMS-UNIVERSITÄT
BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY
Email address: sam.hughes.maths@gmail.com; hughes@math.uni-bonn.de

(S. Machado) DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, SWITZERLAND
Email address: simon.machado@math.ethz.ch